

2013 Egyptian Cottonworm (*Spodoptera littoralis*) Survey Summary

Report Generated:	11/5/2013 10:26
Data Range:	6/3/2013 - 9/3/2013

* Generated by Wyoming Pest Detection Program - www.uwyo.edu/capsweb

COUNTY	S. LITTORALIS TRAPS OR SURVEY SITES		NEW COUNTY RECORD
	NEGATIVE	POSITIVE	
ALBANY	0	0	NO
BIG HORN	5	0	NO
CAMPBELL	0	0	NO
CARBON	0	0	NO
CONVERSE	0	0	NO
CROOK	0	0	NO
FREMONT	1	0	NO
GOSHEN	5	0	NO
HOT SPRINGS	0	0	NO
JOHNSON	0	0	NO
LARAMIE	5	0	NO
LINCOLN	5	0	NO
NATRONA	0	0	NO
NIOBRARA	0	0	NO
PARK	5	0	NO
PLATTE	5	0	NO
SHERIDAN	0	0	NO
SUBLETTE	0	0	NO
SWEETWATER	0	0	NO
TETON	0	0	NO
UINTA	0	0	NO
WASHAKIE	5	0	NO
WESTON	0	0	NO
YELLOWSTONE NATIONAL PARK	0	0	NO
TOTAL	36	0	0

S. littoralis was not found in Wyoming in 2013

36 Traps or Survey Sites were examined for S. littoralis in 8 Wyoming Counties

- 1 Trap or Survey Site was in Apple
- 0 Traps or Survey Sites were in Plum
- 0 Traps or Survey Sites were in Crabapple
- 0 Traps or Survey Sites were in Grape
- 20 Traps or Survey Sites were in Barley
- 0 Traps or Survey Sites were in Oats
- 15 Traps or Survey Sites were in Wheat

36 Traps were examined:

- 33 Traps were placed in Crops
 - 0 Traps were placed in Home Gardens or Yards
 - 0 Traps were placed in Nurseries
 - 1 Trap was placed in an Orchard
 - 2 Traps were placed in Research / Experiment Areas
 - 0 Traps were placed in Vineyards