

Historical Egyptian Cottonworm (*Spodoptera littoralis*) Survey Summary

Report Generated:	1/21/2022 14:07
Data Range:	5/17/2007 - 9/14/2021

* Generated by Wyoming Pest Detection Program - www.uwyo.edu/capsweb

COUNTY	S. LITTORALIS TRAPS OR SURVEY SITES		NEW COUNTY RECORD
	NEGATIVE	POSITIVE	
ALBANY	4	0	NO
BIG HORN	57	0	NO
CAMPBELL	16	0	NO
CARBON	1	0	NO
CONVERSE	1	0	NO
CROOK	16	0	NO
FREMONT	13	0	NO
GOSHEN	76	0	NO
HOT SPRINGS	7	0	NO
JOHNSON	1	0	NO
LARAMIE	69	0	NO
LINCOLN	41	0	NO
NATRONA	2	0	NO
NIOBRARA	1	0	NO
PARK	58	0	NO
PLATTE	69	0	NO
SHERIDAN	7	0	NO
SUBLETTE	1	0	NO
SWEETWATER	1	0	NO
TETON	1	0	NO
UINTA	1	0	NO
WASHAKIE	59	0	NO
WESTON	1	0	NO
YELLOWSTONE NATIONAL PARK	0	0	NO
TOTAL	503	0	0

S. littoralis has not been found in Wyoming

503 Traps or Survey Sites have been examined for S. littoralis in 23 Wyoming Counties

- 16 Traps or Survey Sites have been in Apple
- 2 Traps or Survey Sites have been in Plum
- 3 Traps or Survey Sites have been in Corn
- 10 Traps or Survey Sites have been in Crabapple
- 15 Traps or Survey Sites have been in Grape
- 232 Traps or Survey Sites have been in Barley
- 3 Traps or Survey Sites have been in Oats
- 8 Traps or Survey Sites have been in Spelt
- 214 Traps or Survey Sites have been in Wheat

503 Traps have been examined:

- 436 Traps have been placed in Crops
- 9 Traps have been placed in Home Gardens or Yards
- 18 Traps have been placed in Nurseries
- 1 Trap has been placed in an Orchard
- 26 Traps have been placed in Research / Experiment Areas
- 1 Trap has been placed in a Right-of-Way
- 12 Traps have been placed in Vineyards