

Historical Egyptian Cottonworm (*Spodoptera littoralis*) Survey Summary

Report Generated:	10/10/2016 11:01
Data Range:	5/17/2007 - 8/24/2016

* Generated by Wyoming Pest Detection Program - www.uwyo.edu/capsweb

COUNTY	S. LITTORALIS TRAPS OR SURVEY SITES		NEW COUNTY RECORD
	NEGATIVE	POSITIVE	
ALBANY	4	0	NO
BIG HORN	37	0	NO
CAMPBELL	16	0	NO
CARBON	1	0	NO
CONVERSE	1	0	NO
CROOK	16	0	NO
FREMONT	13	0	NO
GOSHEN	48	0	NO
HOT SPRINGS	7	0	NO
JOHNSON	1	0	NO
LARAMIE	43	0	NO
LINCOLN	21	0	NO
NATRONA	2	0	NO
NIOBRARA	1	0	NO
PARK	38	0	NO
PLATTE	43	0	NO
SHERIDAN	4	0	NO
SUBLETTE	1	0	NO
SWEETWATER	1	0	NO
TETON	1	0	NO
UINTA	1	0	NO
WASHAKIE	39	0	NO
WESTON	1	0	NO
YELLOWSTONE NATIONAL PARK	0	0	NO
TOTAL	340	0	0

S. littoralis has not been found in Wyoming

340 Traps or Survey Sites have been examined for S. littoralis in 23 Wyoming Counties

- 16 Traps or Survey Sites have been in Apple
- 2 Traps or Survey Sites have been in Plum
- 10 Traps or Survey Sites have been in Crabapple
- 12 Traps or Survey Sites have been in Grape
- 154 Traps or Survey Sites have been in Barley
- 1 Trap or Survey Site has been in Oats
- 145 Traps or Survey Sites have been in Wheat

340 Traps have been examined:

- 289 Traps have been placed in Crops
- 9 Traps have been placed in Home Gardens or Yards
- 18 Traps have been placed in Nurseries
- 1 Trap has been placed in an Orchard
- 14 Traps have been placed in Research / Experiment Areas
- 9 Traps have been placed in Vineyards