

**Historical Egyptian Cottonworm (*Spodoptera littoralis*) Survey Summary**

Report Generated:	9/18/2017 11:20
Data Range:	5/17/2007 - 8/25/2017

\* Generated by Wyoming Pest Detection Program - [www.uwyo.edu/capsweb](http://www.uwyo.edu/capsweb)

COUNTY	S. LITTORALIS TRAPS OR SURVEY SITES		NEW COUNTY RECORD
	NEGATIVE	POSITIVE	
ALBANY	4	0	NO
BIG HORN	42	0	NO
CAMPBELL	16	0	NO
CARBON	1	0	NO
CONVERSE	1	0	NO
CROOK	16	0	NO
FREMONT	13	0	NO
GOSHEN	53	0	NO
HOT SPRINGS	7	0	NO
JOHNSON	1	0	NO
LARAMIE	48	0	NO
LINCOLN	26	0	NO
NATRONA	2	0	NO
NIOBRARA	1	0	NO
PARK	43	0	NO
PLATTE	48	0	NO
SHERIDAN	4	0	NO
SUBLETTE	1	0	NO
SWEETWATER	1	0	NO
TETON	1	0	NO
UINTA	1	0	NO
WASHAKIE	44	0	NO
WESTON	1	0	NO
YELLOWSTONE NATIONAL PARK	0	0	NO
<b>TOTAL</b>	<b>375</b>	<b>0</b>	<b>0</b>

**S. littoralis has not been found in Wyoming**

**375 Traps or Survey Sites have been examined for S. littoralis in 23 Wyoming Counties**

- 16 Traps or Survey Sites have been in Apple
- 2 Traps or Survey Sites have been in Plum
- 10 Traps or Survey Sites have been in Crabapple
- 12 Traps or Survey Sites have been in Grape
- 176 Traps or Survey Sites have been in Barley
- 1 Trap or Survey Site has been in Oats
- 158 Traps or Survey Sites have been in Wheat

**375 Traps have been examined:**

- 322 Traps have been placed in Crops
  - 9 Traps have been placed in Home Gardens or Yards
  - 18 Traps have been placed in Nurseries
  - 1 Trap has been placed in an Orchard
- 16 Traps have been placed in Research / Experiment Areas
  - 9 Traps have been placed in Vineyards