


- 1 This RAAT method has been adopted in a 2EE label for most western states.
- 2 Increasing the volume of water above the 1:1 ratio with carbaryl will not improve, and may reduce, efficacy.
- 3 Facilitating droplet dispersal by using Micronaire nozzles or turning conventional nozzles angled 45 degrees into the wind may enhance efficacy. In addition, wider treated swaths (>100 foot) may be used. While it may be possible to proportionately expand untreated buffers, this approach has not been tested. Therefore, the most prudent tactic at this time would be to maintain the fixed widths of the untreated swaths.
- 4 Experimental applications have demonstrated that RAATs provide more than 90 percent mortality when ground temperatures exceed air temperatures by as much as 15 degrees Fahrenheit. Extensive research in Australia has demonstrated that when ground temperatures exceed air temperatures, thermals can limit insecticide deposition of aerial applications unless there are winds of 4 to 10 miles per hour (ideally, perpendicular to the swaths).
- 5 Using canola or corn oil, rather than crop oil, as a carrier may significantly improve the effectiveness of RAAT applications. Canola and corn oils are effective attractants and feeding stimulants for many rangeland grasshopper species.
- 6 Small-scale trials under optimal conditions indicate that 80 to 85 percent mortality also can be achieved using 100-foot swaths with the following rates and coverages: 5 fluid ounces per acre with 33-foot untreated swaths, 6 ounces with 50-foot untreated swaths, and 7 ounces with 100-foot untreated swaths.

For more information, visit:  
[www.wygisc.uwyo.edu/grasshopper/](http://www.wygisc.uwyo.edu/grasshopper/)  
 or contact

Professor Jeffrey A. Lockwood  
 Association for Applied Acridology International  
 Department of Renewable Resources  
 University of Wyoming College of Agriculture  
 P.O. Box 3354  
 Laramie, WY 82071-3354  
 (307)766-4260, lockwood@uwyo.edu

For more information on RAAT, visit [www.wygisc.uwyo.edu/grasshopper/](http://www.wygisc.uwyo.edu/grasshopper/) (UWyo link)  
 Dr. Jeffrey A. Lockwood, 1000 E. University Ave., Dept. of Renewable Resources, U.S. Department of Agriculture, APIS, 1400 N. Lincoln Ave., Laramie, WY 82071-3354, lockwood@uwyo.edu  
 Dr. Jeffrey A. Lockwood, 1000 E. University Ave., Dept. of Renewable Resources, U.S. Department of Agriculture, APIS, 1400 N. Lincoln Ave., Laramie, WY 82071-3354, lockwood@uwyo.edu

### A schematic of an aerial RAAT application with 50 percent coverage.



### Exceptions to the “Rules”

Higher rates or coverages may be needed if: 1) treatments are applied to late instar nymphs (if using diflubenzuron), 2) ground temperatures exceed air temperatures (especially if using malathion), 3) grasshopper densities are extreme, 4) forage cover is tall or dense, or 5) terrain is rough. In all cases, grasshopper management software (CARMA<sup>1</sup> or HOPPER<sup>2</sup>) should be used to assess a program. Apply insecticides in accordance with label directions and established guidelines for buffers around water, bees, and human habitations.

<sup>1</sup> Available at [www.wygisc.uwyo.edu/grasshopper/carma.htm](http://www.wygisc.uwyo.edu/grasshopper/carma.htm)  
<sup>2</sup> Available from your USDA APHIS State Plant Health Director


**UNIVERSITY OF WYOMING**  
 1000 E. UNIVERSITY AVENUE  
 LARAMIE, WY 82071-3354  
 (307) 766-4260  
[www.uwyo.edu](http://www.uwyo.edu)

**AERIAL-RAAT STRATEGY**

Do more with less,  
 using  
**Reduced Agent and Area Treatments (RAATs)**

*Refining*

## What are RAATs?

RAAT is a method of integrated pest management (IPM) for rangeland grasshoppers in which the rate of insecticide is reduced from traditional levels and untreated swaths (refuges) are alternated with treated swaths.

RAATs work through *selective mortality*, meaning grasshoppers are killed directly in treated swaths and as they move out of untreated swaths, and *biological control*, which allows predators and parasites preserved in untreated swaths to suppress grasshoppers. This IPM approach can reduce the cost of control and the amount of insecticide used by more than 50 percent.

## Why use RAATs?

### Economics

Inflation and changes in governmental cost-share programs have tripled the cost of rangeland grasshopper management for ranchers compared with the control cost during the last major outbreak in 1987. Most states no longer subsidize grasshopper control programs; therefore, the cost of traditional tactics usually will exceed the benefits.

### Environment

Less insecticide in the environment lowers the risk to native species (including fish and wildlife), water quality, and humans. The untreated swaths provide a refuge for organisms with lower mobility than grasshoppers, and even those organisms that move into the treated swaths will be largely unaffected unless they feed on the foliage.

## What to expect from RAATs

### Efficacy

This method normally will result in 80 to 95 percent control, which is approximately 5 to 15 percent lower mortality than with a standard (high rate, blanket coverage) treatment. Leaving low densities of grasshoppers after RAATs does not necessarily result in a subsequent outbreak (see *Economic Benefits*).

### Economics

Using RAATs will reduce costs by approximately 50 to 60 percent, depending upon the agent and swath width. For example, if a standard insecticide application costs \$2.50 per acre, the equivalent RAAT program should cost approximately \$1.15 per acre prevented. In some cases, costs are reduced by two-thirds. It should be noted that the greatest economical benefits derive from increased swath spacing, which also decreases the costs for both purchasing insecticides and application.

### Environment

Labels mean 50 to 75 percent less insecticide is applied to our rangelands for grasshopper control. The untreated swaths harbor species essential to rangeland ecosystems, including biocontrol agents of grasshoppers and weeds. Low densities of surviving grasshoppers allow predators and parasites in the untreated refuges to recolonize and thereby reestablish natural regulation of grasshopper populations. For these reasons, RAAT programs

