## Homework 4 (MATH 5490-01)Name (Print):Due date: Thursday, April 11, 2013

1. Consider the following model for the PDF f(v, t) of molecular velocities v in one direction:

$$\frac{\partial f(\mathbf{v}, \mathbf{t})}{\partial \mathbf{t}} = \frac{\partial}{\partial \mathbf{v}} \left[ \frac{\mathbf{v} - \mathbf{V}}{\tau_{m}} f(\mathbf{v}, \mathbf{t}) \right] + \frac{2e}{3\tau_{m}} \frac{\partial^{2} f(\mathbf{v}, \mathbf{t})}{\partial v^{2}}.$$
(1)

Here,  $\tau_m$  is the constant characteristic time scale of molecular velocity fluctuations, and V and e are constant model parameters.

a) Calculate the stationary PDF f(v) (which is independent of time t) by integration of equation (1).Hint: The first integration constant can be determined by the constraint that both

f(v) and  $\partial f(v)/\partial v$  have to be zero if  $v \to \infty$ . The second integration constant can be determined by the normalization constraint for f(v).

- b) Explain the meaning of V and e by relating these variables to the mean and variance of f(v). Explain why the stationary PDF f(v) does not depend on  $\tau_m$ .
- 2. The development of a certain population in time t is described by the following equation for the population PDF f(p, t):

$$\frac{\partial f(\mathbf{p},t)}{\partial t} = \frac{\partial}{\partial \mathbf{p}} \left[ \frac{\mathbf{p} - \mathbf{C}}{\tau} f(\mathbf{p},t) \right] + \frac{\partial^2 \mathbf{D} f(\mathbf{p},t)}{\partial \mathbf{p}^2}.$$
(2)

Here,  $\tau$  is the constant characteristic time scale of population fluctuations, and C and D are constant model parameters.

- a) Use equation (2) to find the mean <P> and variance  $<\tilde{P}^2>$  as functions of t.
- b) Find the conditional PDF f(p, t | p', 0) such that its parameters α and β are given as explicit functions of t.

Hint: The best way to address this question is to consider the equations for  $\alpha - \langle P \rangle$ and  $\beta - \langle \tilde{P}^2 \rangle$ .

- c) Write the PDF f(p, t) in dependence on the conditional PDF f(p, t | p', 0) and any initial PDF f(p', 0). What is the PDF f(p, t) if  $t \rightarrow \infty$ ?
- d) Calculate the asymptotic  $(t \rightarrow \infty)$  PDF f(p, t) for the case that D  $\rightarrow$  0. Explain the meaning of your result.