

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
FOR ANIMAL TREATMENT ONLY

# ivomec<sup>®</sup>

(ivermectin)

## Pour-On for Cattle

Active Constituent: IVERMECTIN 5 mg/mL

For the treatment and control of ivermectin-sensitive gastrointestinal roundworms (including inhibited *Ostertagia ostertagi*), lungworm (*Dictyocaulus viviparus*), eyeworm (*Thelazia* spp.), sucking and biting lice and sarcoptic and chorioptic mange mites, buffalo fly (*Haematobia irritans exigua*) and cattle tick (*Boophilus microplus*) in cattle, and for the treatment and control of gastrointestinal roundworms and lungworm of red deer.



**READ ENCLOSED LEAFLET BEFORE USING THIS PRODUCT**

Dose 1mL per 10kg liveweight  
For external use only



Merial Australia Pty Limited  
A.C.N. 071 187 285  
Level 6, 79 George Street  
Parramatta NSW 2150

For product information and for emergency advice call the 24-hour toll-free Merial technical hotline 1800-808-691.

NRA Approval No.: 45721/0999

IVOMEc and the cattle head logo are registered trademarks of Merial.  
© 1999 Merial.

**20 Litre**

**Product No.: H 41364X (AU)**



49205/021199C

2e PROEF  
PDF: 10-11-1999

**ivomec**<sup>®</sup> Pour-On for Cattle  
(ivermectin)

Product No.:H 41364X (AU)

20 Litre

**INDICATIONS:**

For the treatment and control of ivermectin-sensitive gastrointestinal roundworms (including inhibited *Ostertagia ostertagi*), lungworm (*Dictyocaulus viviparus*), eyeworm (*Thelazia* spp.), sucking and biting lice and sarcoptic and chorioptic mange mites, buffalo fly (*Haematobia irritans exigua*) and cattle tick (*Boophilus microplus*) in cattle, and for the treatment and control of gastrointestinal roundworms and lungworm of red deer.

**PERSISTENT ACTIVITY:**

'IVOMEC' Pour-On for Cattle applied at the recommended dose of 1mL per 10kg of liveweight controls infections of *Trichostrongylus axei* (stomach hair worm) and *Cooperia* spp. (small intestinal worm) for up to 14 days after treatment; *Ostertagia* spp. (small brown stomach worm) and *Oesophagostomum radiatum* (nodule worm) for up to 21 days after treatment; *Dictyocaulus viviparus* (lungworm, in deer as well as cattle) for up to 28 days after treatment; *Haematobia irritans exigua* (buffalo fly) for up to 21 days after treatment.

**DOSAGE AND ADMINISTRATION:**

'IVOMEC' Pour-On for Cattle is for external application to cattle and red deer only. Apply along the topline in a narrow strip extending from the withers to the tailhead. The dose rate is 1mL for each 10kg of liveweight.

It is generally recommended that no topical preparations should be used during rainfall, or when cattle or deer are very wet. However, the effect of 'IVOMEC' Pour-On for Cattle is not adversely affected if cattle hide is wet or if rain occurs shortly after application.

**WITHHOLDING PERIODS:**

**MEAT: DO NOT APPLY TO CATTLE LESS THAN 42 DAYS BEFORE SLAUGHTER FOR HUMAN CONSUMPTION.**

**DO NOT APPLY TO DEER LESS THAN 35 DAYS BEFORE SLAUGHTER FOR HUMAN CONSUMPTION.**

**MILK: WHEN MILK OR MILK PRODUCTS ARE TO BE USED FOR HUMAN CONSUMPTION DO NOT ADMINISTER TO COWS WITHIN 28 DAYS PRIOR TO CALVING OR DURING LACTATION.**

**PRECAUTIONS:**

Use only in well-ventilated areas or outdoors.

Close container when not in use.

Cloudiness may result when 'IVOMEC' Pour-On for Cattle is stored at temperatures below 0° C. Allowing to warm at room temperature will restore the normal appearance without affecting efficacy.

The antiparasitic activity of ivermectin will be impaired if the formulation is applied to areas of skin with mange scabs or lesions, or with dermatoses or adherent materials eg. caked mud or manure.

**This product is for application to the skin surface only, do not give orally or parenterally.**

'IVOMEC' Pour-On for Cattle is not recommended for use in species other than cattle and red deer.

Treatment should not be repeated in less than 42 days.

**NOTE TO USER:**

The colour of 'IVOMEC' Pour-On for Cattle fades when exposed to light, and depending on the light intensity, fading may occur in less than 30 minutes. This rapid loss of colour does not reflect loss of potency of ivermectin. However, prolonged exposure (i.e. weeks) to light can also result in a gradual decline of ivermectin potency in the formulation.



49206/021199C

2e PROEF  
PDF: 10-11-1999

**ivomec**<sup>®</sup> Pour-On for Cattle  
(ivermectin)

**Product No.:H 41364X (AU)**  
**20 Litre**

**DISPOSAL:**

Break, crush or puncture and bury empty containers at a local authority landfill. If not available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt. Do not dispose of undiluted chemicals on site.

**PROTECTION OF FISH, CRUSTACEANS AND THE ENVIRONMENT**

Studies indicate that when ivermectin comes in contact with the soil, it readily and tightly binds to the soil and becomes inactive over time. Containers and any residual contents should be disposed of safely in an approved landfill as free ivermectin may adversely affect fish and certain water-borne organisms.

**DO NOT CONTAMINATE DAMS, RIVERS OR STREAMS.**

**SAFETY DIRECTIONS:**

**Poisonous if swallowed.**

**Avoid contact with eyes, skin and clothing.**

**FIRST AID:**

**If poisoning occurs contact a doctor or Poisons Information Centre, phone 131126.**

**For further information, refer to the Material Safety Data Sheet.**

**DANGER**

**GIVES OFF HIGHLY FLAMMABLE VAPOUR.**

**KEEP AWAY FROM HEAT, SPARKS, OPEN FLAME AND OTHER SOURCES OF IGNITION.**

**KEEP CLOSED WHEN NOT IN USE.**

**STORAGE:**

Store below 30°C (Room Temperature).

Store bottle in carton to protect from light.

Batch No.:

Expiry Date:



UN 1219 iso-PROPANOL



49207/021199C