

Boehringer
Ingelheim

Revision 7
Operator hal
Date 28Oct13
Art # 384109-00
Item # 81647755
Product Cydectin Injectable

Type/Size Label/500 mL
Template # 156
Dimensions 3-1/4" x 7"
Corner Radius125"
PDF 28Oct13 hal
Rev Level 00

BLACK

PMS 8243

PMS 877

PMS 8825

UNVARNISHED
AREA 1" X 2"

NDC 0010-3841-02

CYDECTIN®

moxidectin

Injectable Solution for Beef and Nonlactating Dairy Cattle

Antiparasitic

For Treatment of Infections and Infestations Due to Internal and External Parasites of Cattle.

DOSE: The recommended rate of administration for CYDECTIN Injectable is 1 mL for each 110 lb (50 kg) body weight to provide 0.2 mg moxidectin/2.2 lb (0.2 mg/kg) body weight. Be careful not to overdose animals; estimate animal's weight as closely as possible or weigh animals individually.

PRECAUTIONS: This product is intended for subcutaneous use in cattle only and should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result.

STORAGE: Store product at or below 77°F (25°C). Protect from light.

DISPOSAL: Do not contaminate water by direct application or by improper disposal of drug container. Dispose of containers in an approved landfill or by incineration.

Read accompanying package insert carefully before use.

Restricted Drug (CA) - Use Only As Directed

Sterile

Contains 10 mg moxidectin/mL (5,000 mg/500 mL)

Not for use in female dairy cattle
20 months of age or older
(including dry dairy cows),
veal calves, and calves less
than 8 weeks of age.

Contents **500 mL**



NADA 141-220, Approved by FDA

HUMAN WARNINGS: Not For Use in Humans. Keep this and all drugs out of the reach of children.

To report adverse reactions attributable to exposure to this product, call 1-866-638-2226.

RESIDUE WARNINGS: Cattle must not be slaughtered for human consumption within 21 days of treatment. This drug product is not approved for use in female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows. A withdrawal period has not been established for preruminating calves. Do not use in calves to be processed for veal.

ANIMAL SAFETY WARNINGS: Do not use in sick, debilitated, or underweight animals. In foreign countries there have been reports of adverse effects, including death. This product should not be used in calves less than 8 weeks of age because safety testing has not been done in the U.S. in calves less than 8 weeks of age.

U.S. Patent No. 5,965,603 - used under license.

© 2013 Boehringer Ingelheim Vetmedica, Inc.
All Rights Reserved.

Manufactured for:
Boehringer Ingelheim Vetmedica, Inc.
St. Joseph, MO 64506 U.S.A.

B06K3

Lot No.:
Exp. Date:

384109-00
81449783
81647755, R.0