

# **Wetland Profile and Condition Assessment of the Little Snake River, Wyoming**

FINAL REPORT

September 26th, 2018



Lindsey Washkoviak, Teresa M. Tibbets, and George Jones

Wyoming Natural Diversity Database, University of Wyoming, 1000 East University Avenue,  
Department 3381, Laramie, WY 82071

## **Wetland Profile and Condition Assessment of the Little Snake River, Wyoming**

**Prepared for:**

U.S. Environmental Protection Agency, Region 8  
1595 Wynkoop Street  
Denver, CO 80202

EPA Assistance ID No. CD - 96805101

**Prepared by:**

**Lindsey Washkoviak, George Jones, and Teresa M. Tibbets**  
**Wyoming Natural Diversity Database**  
University of Wyoming  
Department 3381, 1000 East University Avenue  
Laramie, WY 82071

This document should be cited as follows:

L. Washkoviak, G. Jones, and Tibbets, T. M. 2018. Wetland Profile and Condition Assessment of the Little Snake River, Wyoming. Prepared for the U.S. Environmental Protection Agency by the Wyoming Natural Diversity Database, University of Wyoming, Laramie, Wyoming. 41 pp. plus appendices.

Cover photograph: Scott Copeland

# TABLE OF CONTENTS

LIST OF TABLES .....	iii
LIST OF FIGURES .....	iv
EXECUTIVE SUMMARY .....	v
ACKNOWLEDGEMENTS .....	vii
1.0 INTRODUCTION .....	1
1.1 Wyoming’s strategy for wetland assessments.....	1
1.2 Project Background .....	1
2.0 STUDY AREA .....	2
2.1 Land Use .....	3
2.2 Topography .....	3
2.3 Hydrology.....	3
3.0 LANDSCAPE PROFILE OF THE LITTLE SNAKE RIVER BASIN .....	4
3.1 Wetland Resource Description.....	5
3.1.1 <i>Wetland Class</i> .....	5
3.1.2 <i>Water Regime</i> .....	5
3.1.3 <i>Special modifiers describing wetlands</i> .....	7
3.2 Land Ownership/Management .....	8
4.0 WETLAND CONDITION ASSESSMENT .....	9
4.1 Ecological Integrity Assessment Framework.....	9
4.2 Survey Design and Evaluation of Sample Sites .....	10
4.2.1 <i>Wetland Definitions for Target Population</i> .....	10
4.2.2 <i>Sample Frame and Classification</i> .....	10
4.2.3 <i>Sample Size and Selection Criteria for Site Evaluation</i> .....	11
4.3 Field Methods.....	12
4.3.1 <i>Wetland Assessment Area (AA)</i> .....	12
4.3.2 <i>Ecological Integrity Assessment of Wetland Sites</i> .....	13
4.3.3 <i>Landscape Hydrology Metric (LHM)</i> .....	13
4.3.4 <i>Vegetation Assessments</i> .....	14
4.3.5 <i>Soils</i> .....	15

4.3.6 <i>Surface Water Characterization</i> .....	15
4.4 Data Analysis .....	15
4.4.1 <i>Data Management</i> .....	15
4.4.2 <i>Ecological Integrity Assessment</i> .....	15
4.4.3 <i>Floristic Quality Assessment (FQA)</i> .....	16
5.0 WETLAND CONDITION ASSESSMENT RESULTS .....	17
5.1 Characteristics of Sampled Wetlands .....	17
5.1.1 <i>Characteristics of Sampled Wetland Subgroups</i> .....	18
5.2 Characterization of Wetland Vegetation .....	20
5.2.1 <i>Species Diversity of Wetland Vegetation</i> .....	20
5.2.2 <i>Floristic Quality Assessment</i> .....	21
5.2.3 <i>Plant Species of Conservation Concern</i> .....	22
5.3 Wildlife Observations .....	24
5.4 Wetland Condition Assessment .....	25
5.4.1 <i>Ecological Integrity Assessment of Sampled Wetlands</i> .....	25
5.4.2 <i>Estimate of Wetland Condition for targeted wetlands in the LSRB</i> .....	27
5.4.3 <i>Landscape Hydrology Metric</i> .....	28
5.4.4 <i>Indicators of disturbance</i> .....	28
6.0 LITTLE SNAKE FOCAL AREAS .....	29
6.1 The Muddy Creek Wetlands Project .....	30
6.2. Upper Basin Vs Floodplain .....	31
6.2.1 <i>Wetland Types</i> .....	31
6.2.2 <i>Ecological Condition</i> .....	32
6.2.3 <i>Indicators of Disturbance</i> .....	32
6.2.4 <i>Hydrologic Alteration</i> .....	33
7.0 DISCUSSION .....	35
8.0 CONCLUSION .....	35
LITERATURE CITED .....	37

## LIST OF TABLES

Table 1. Surface area of wetlands based on NWI classifications in the LSRB. ....	5
Table 2. Percent of wetlands with a specific hydrologic regime in the LSRB. ....	6
Table 3. Area of wetland in acres classified according to NWI modifiers in the LSRB. ....	7
Table 4. Areas of irrigated wetlands in acres based on NWI classifications in the LSRB. ....	8
Table 5. Land ownership/management of wetlands by area in the LSRB. ....	9
Table 6. Target wetland subgroups classified by Cowardin, Hydrogeomorphic (HGM), and Ecological Systems used in the LSRB. ....	11
Table 7. EIA attributes and metrics used for wetland assessments in the LSRB. ....	13
Table 8. Landscape Hydrology Metric scoring criteria. ....	14
Table 9. Number of wetland sites in each target wetland subgroup evaluated, sampled, and rejected during the study by wetland subgroup. ....	17
Table 10. Reasons for rejection of wetlands in the LSRB during the study. ....	17
Table 11. Ten most common plant species documented at sampled wetland in the LSRB. ....	21
Table 12. Frequencies of native and non-native species encountered at the sites sampled in the LSRB. ....	21
Table 13. Floristic quality assessment indices calculated for wetlands in the LSRB. ....	22
Table 14. Plant species of concern identified at sampled wetlands within the LSRB. ....	23
Table 15. Wildlife observations made during wetland sampling effort in the LSRB. ....	24
Table 16. Ranks for each EIA attribute by wetland subgroup for the LSRB. ....	26
Table 17. Population estimate of EIA ranks for wetlands in the LSRB. Observed = percent of sampled sites within each rank; Estimate = percent of wetland number extrapolated using 67 wetlands from the sample frame. ....	27
Table 18. Number of wetland sites sampled by landowner/manager in the Upper Basin and Floodplain focal areas. ....	31
Table 19. Number of wetland sites sampled by wetland classification in the Upper Basin and Floodplain focal areas. ....	32
Table 20. Number of wetland sites sampled by ecological condition score in the Upper Basin and Floodplain focal areas. ....	32
Table 21. Number of wetland sites sampled with disturbance indicators present in the Upper Basin and Floodplain focal areas. ....	33
Table 22. Number of wetland sites sampled by LHM Classification in the Upper Basin and Floodplain focal areas. ....	34

**LIST OF FIGURES**

Figure 1. The Little Snake River Basin study area ..... 2

Figure 2. 8 USGS quads that will have updated wetland mapping available in 2019..... 4

Figure 3. Surface area (acres) of wetlands classified according to NWI water regime in the LSRB..... 6

Figure 4. Spatial distribution of land ownership/management and wetlands within the LSRB study area. .... 8

Figure 5. Description of land ownership/management (acres) of the study area in the LSRB..... 9

Figure 6. Percentage of points selected in the original survey design versus sites sampled by landowner/manager ..... 18

Figure 7. Riverine shrubland wetlands in the LSRB. .... 19

Figure 8. Freshwater emergent marsh wetlands in the LSRB. .... 19

Figure 9. Wet meadow wetlands in the LSRB..... 20

Figure 10. Sixty-seven percent of the wetlands in the LSRB received an EIA score of B or above. EIA scores were lowest for emergent marshes, indicating 50% of these wetlands surveyed received a C or below, indicating moderate to significant departure from reference..... 25

Figure 11. Cumulative distribution function of wetland EIA scores with 95% CI shown. Graph is the cumulative proportion of wetlands (y-axis) with EIA scores at or below values on the x axis. .... 27

Figure 12. Proportion of total wetland sites based on categories for the Landscape Hydrology Metric..... 28

Figure 13. Potential indicators of disturbance observed across all wetlands in the Little Snake River Basin..... 29

Figure 14. Focal areas within the LSRB showing landownership/management ..... 30

Figure 15. Spatial distribution of wetland acres, irrigated acres, and wetlands acres overlapping with irrigated acres in the Floodplain focal area..... 34

## EXECUTIVE SUMMARY

This report presents the results of a landscape profile and basin-scale evaluation of the ecological condition of wetlands in the Little Snake River Basin (LSRB) of south-central Wyoming. This project is part of a state-wide effort to better understand the extent and ecological condition of wetlands in Wyoming. The objectives of this document are to: [1] describe the landscape profile of wetlands within the LSRB study area and [2] report results from a field-based assessment of wetland condition in the LSRB conducted in 2016.

We developed a multi-level approach to create a landscape profile and estimate condition of wetlands within the LSRB study area. We produced a wetland landscape profile using digital wetland mapping data to describe wetland resources. We assessed basin-scale wetland condition using Ecological Integrity Assessment (EIA) methods supplemented by measurements of anthropogenic and hydrologic disturbance, baseline characteristics of wetland vegetation, and hydrologic alteration. EIA field protocols were used to survey 68 randomly-selected and 2 intentionally-selected sites in June-September 2016. Surveys targeted three wetland subgroups: 1) Wet meadows; 2) Emergent marshes; and 3) Riverine Shrubland.

Based on digital mapping, wetlands within the study area totaled 11,636 acres, or 4% of the LSRB study area. The landscape profile results show the importance of understanding linkages between land use, irrigation practices and wetlands in the LSRB. Over 85% of wetlands in the study area are privately-owned and wetlands comprise approximately half of the irrigated landscape. Over 65% of freshwater emergent wetlands, the most common type, are mapped as irrigated. Coordination with private landowners is essential to maintain the ecological integrity of wetland resources throughout the LSRB.

Our study found that all ecological subgroupings were dominated by B-ranked (slight deviation from reference condition) wetlands, meaning there was evidence of low levels of disturbance and a slight deviation from reference condition. We estimate 7% of wetlands were A-ranked (no deviation from reference condition), 60% B-ranked (moderate deviation from reference condition), 24% C-ranked (moderate impact) and 9% D-ranked (significant deviation from reference condition). Riverine shrublands had the highest proportion of A-ranked sites and no D-ranked sites, indicating overall lower disturbance relative to other wetland types. Wet meadows and emergent marshes scored a C or below for 30% and 50% of sites, respectively.

We collected data documenting potential disturbances that may influence wetland condition. The most widespread disturbances identified in our study were presence of non-native species, roads, grazing by domestic and native herbivores and modified hydrology from irrigation infrastructure. Almost all (97%) sites sampled had non-native species present. We developed a Landscape Hydrology Metric (LHM) that identified altered hydrology at 47% of sampled wetlands. Although irrigation and related agricultural activities are generally considered

disturbance factors, water availability of many wetlands can be enhanced by these anthropogenic activities, especially in arid regions like the LSRB.

After sampling and reviewing condition assessment results, we found that the LSRB study area divided into three distinct areas (the Upper Basin, the Floodplain, and the Muddy Creek Wetlands Project (MCWP) focal areas) that had different landowner/management patterns, land uses, wetland types, and potential patterns of disturbance. All sampled wetlands in the Floodplain were affected by altered hydrology and lower ecological condition (71% of C-ranked wetlands and all D-ranked wetlands). Indicators of disturbance were associated with agriculture and development. In contrast, all A-ranked wetlands and most (88%) B-ranked wetlands were located in the Upper Basin where 80% of sampled wetlands lacked hydrologic alterations. Indicators of disturbance were related to grazing and soil degradation by native ungulates and livestock. Two wetlands were sampled at the MCWP and received an ecological condition ranking of a C. Wetlands at the MCWP are unique in the basin because they were created as part of a restoration effort to provide habitat for migratory waterfowl and shorebirds, and present difficulties when assessing ecological condition.



## ACKNOWLEDGEMENTS

This project was funded by a Wetland Program Development Grant (#CD - 96805101) from the U.S. Environmental Protection Agency Region 8 and the Wyoming Chapter of The Nature Conservancy. The framework for this study was informed by the State Wildlife Wetlands Conservation Strategy (Wyoming Joint Ventures Steering Committee 2010), the State Wildlife Action Plan (Wyoming Game and Fish Department 2010), and the Wyoming Bird Conservation Habitat Partnership (WBCHP) including representatives from Wyoming Game and Fish Department (WGFD), Wyoming Department of Environmental Quality (DEQ), USFWS (United States Fish and Wildlife Service), The Nature Conservancy (TNC), Ducks Unlimited (DU), Intermountain West Joint Ventures (IMWJV), Wyoming Audubon, and Rocky Mountain Bird Observatory (RMBO).

We would like to thank Larry Hicks with the Little Snake Conservation District for his support of this project and his help with establishing landowner contacts and community outreach. Thanks to Chad Rieger at Wyoming DEQ for providing guidance and support for the study proposal. We would also like to thank Holly Copeland and Amy Pocewitz from the Wyoming Chapter of The Nature Conservancy for contributing matching funds and support for the project. We extend our gratitude to Mark Andersen at WYNDD for assistance with survey design. We thank our field technician, Chris Nieters, for his hard work collecting and entering data.

This study would not be possible without the public land managers and private landowners that granted access to wetlands on their lands. We extend our gratitude to landowners for their support of this project.

## **1.0 INTRODUCTION**

### **1.1 Wyoming's strategy for wetland assessments**

Freshwater wetland ecosystems are highly diverse, productive habitats, which provide vital ecosystem services (Costanza et al. 1997). Dahl (1990) estimates 38% of wetlands that existed in Wyoming prior to European settlement were lost between 1780 and the mid-1980s. Wetlands remain highly threatened and are subjected to pressures from agricultural, residential, and energy development in Wyoming (Copeland et al. 2010, Pocewicz et al. 2014). Recent studies identify Wyoming's wetlands as one of the habitat types most vulnerable to impacts of future habitat alteration and climate change (Copeland et al. 2010a, Pocewicz et al. 2014).

In light of the changes and threats to wetland ecosystems, we undertook this project as part of a state-wide effort to fill gaps in our understanding of the current extent and ecological condition of wetlands in Wyoming. Recent studies of wetlands in the Intermountain West, including Wyoming, (Lemly and Gilligan 2012, Newlon et al. 2013, Tibbets et al. 2015, 2016a, 2016b) have utilized landscape profiles and rapid assessment methods (RAMs) to draw conclusions regarding wetland resources. Landscape profiles primarily use digital spatial information to quantify the distribution of resources, such as wetland types or area, and to develop conservation goals at a landscape scale (Gwin et al. 1999). RAMs assess the condition of wetlands based on field surveys that measure abiotic and biotic indicators of ecological function and indicators of disturbance that have the potential to negatively affect wetlands. Together, landscape profiles and RAMs are used to establish baseline wetland profiles that include ecological condition, assessment of cumulative impacts, and information useful to prioritize protection and restoration efforts (Gwin et al. 1999).

### **1.2 Project Background**

This report summarizes results of the first basin-wide assessment of wetlands in the Little Snake River Basin (LSRB) of south-central Wyoming. This project was the fifth basin-scale wetland condition assessment conducted in Wyoming, and builds upon previous studies completed within the Laramie Plains Basin (Tibbets et al. 2016), the Goshen Hole Basin (Tibbets et al. 2016b), the Upper Green River Basin (Tibbets et al. 2015), and a statewide landscape scale assessment (Copeland et al. 2010). A report summarizing results of a wetland condition assessment for the Great Divide Basin was completed concurrently as part of this project (Washkoviak et al. 2018)

The need for general information about wetlands in the LSRB is well recognized. This project complements and amplifies the statewide wetland overview completed in 2010 (Copeland et al. 2010), in which the Little Snake wetland complex area was one of nine wetland complexes identified as a statewide-priority area. The LSRB is listed in the Wyoming Wetlands Conservation Strategy (Wyoming Joint Ventures Steering Committee 2010) as primary-focus wetland complex because of unique ecological values and exceptionally high potential for conservation projects. Finally, the Little Snake area lies within the Little Snake/Upper North


Platte Conservation Focus Area of the U.S. Fish and Wildlife Service’s Wyoming Partners for Fish and Wildlife Program (U.S. Fish and Wildlife Service 2012).

The objectives of this document are to: [1] describe the landscape profile of wetlands within the LSRB study area and [2] report results from a field-based assessment of wetland condition in the LSRB conducted in 2016.

## 2.0 STUDY AREA

The LSRB covers 267,098 acres in extreme south-central Wyoming (Figure 1). The study area includes the Little Snake River floodplain and its main tributaries of Cow Creek, Wild Cow Creek, Big Sandstone Creek, and Little Savery Creek.

The study area also includes the Muddy Creek Wetlands Project, the largest constructed wetland complex in Wyoming (WBHCP 2014), which covers 5,000 acres of private and public lands and includes over 2,500 wetland acres located along 6 miles of Muddy Creek and Red Wash Draw near Dad, WY. Numerous constructed impoundments and natural basins occur along the riparian corridor of Muddy Creek to create a complex of emergent marshes and wet meadows intermixed with sage-brush dominated uplands that support hundreds of species of waterfowl and shorebirds during breeding and migration from both the Pacific and Central flyways (Meade 2010).



**Figure 1.** The Little Snake River Basin study area

## **2.1 Land Use**

Land uses in the LSRB are predominantly agriculture-based. This includes sheep and cattle grazing, small scale farming, and cultivated hay crops (WBHCP 2014). Approximately 5% (13,374 acres) of the total basin area is irrigated (Section 3.1.3, Table 4). Over 80% of irrigated lands in the study area (10,907 acres) are located along the southern portion of the Little Snake River floodplain where flood irrigation is used for hay production and cattle grazing.

## **2.2 Topography**

The LSRB is located along the west side of the Sierra Madre and ranges from 6,079 (1853m) to 9,350 ft (2580m) in elevation. The area includes a wide diversity of habitats, from aspen glades, mixed mountain shrubs and sagebrush steppe at higher elevations, to riparian galleries of cottonwood and willow intermixed with herbaceous wetlands at lower elevations (Copeland et al. 2010). The dominant ecological system is Inter-Mountain Basin Montane Sagebrush Steppe (Comer et al. 2003).

## **2.3 Hydrology**


Historically, wetlands in the LSRB consisted of seeps, springs, oxbows and other wetlands associated with the riparian corridor, and to a lesser extent with temporary and seasonal playa wetlands fed mainly by precipitation (WBHCP 2014). Stream flow in the Little Snake River is generally driven by snowpack that accumulates in the mountains and is stored in the High Savery Reservoir during spring runoff for release during the irrigation season. Annual stream flow is naturally highly variable and difficult to predict and is influenced by the High Savery Dam (22,433 acre-foot capacity) and multiple irrigation projects along the Little Snake River (WBHCP 2014).

Water is diverted from the Little Snake River and its tributaries into delivery ditches and canals that convey it by gravity to irrigated fields. The application of irrigation water over time can augment or enlarge historical wetlands or create new wetlands by altering natural hydrology, soil characteristics (e.g., color, redox features, and salt content), and vegetation (U.S. Army Corps of Engineers 2008). Wetlands associated with irrigation in the LSRB include margins of storage reservoirs, seeps along canals and ditches, natural or constructed basins to capture return flows from flood irrigated fields and pastures or overlap with flood irrigation. Irrigation also augments stream flow by contributing late season flows to streams which were historically dry by midsummer, allowing associated wetlands to retain water longer in the season (WBHCP 2014). Overall, these alterations to the hydrology in the LSRB have likely both removed and created wetland area in the basin.

### 3.0 LANDSCAPE PROFILE OF THE LITTLE SNAKE RIVER BASIN

A landscape profile was created using digital wetland mapping data compiled from the U.S. Fish and Wildlife Service (USFWS) National Wetland Inventory (NWI) and additional data layers describing irrigated lands and land ownership within the LSRB study area. The landscape profile describes wetlands within the study area based on codes and modifiers defined by c. by the following attributes: wetland class; water regime; extent of wetlands modified/irrigated (Wyoming Wildlife Consultants 2007); and land management/ownership (Bureau of Land Management 2010). The landscape profile identifies wetlands according to categories

Note: NWI mapping in 8 USGS Quads along the Little Snake River and Muddy Creek Wetlands Project is being updated by St. Mary's University of Minnesota and will be available in 2019 (Figure 2). This update will result in more accurate wetland boundaries. Polygons are being attributed with the Landscape, Landform, Waterbody, Water flow path (LLWW) classification developed by USFWS (Tiner 2003) which can be combined with Cowardin et al. (1979) to estimate functional potential for all wetlands and riparian areas in the LSRB (GeoSpatial Services Saint Mary's University of Minnesota 2018a). This updated NWI mapping will be submitted to the U.S. Fish and Wildlife Service to be added into the national database.



**Figure 2.** 8 USGS quads that will have updated wetland mapping available in 2019.

The landscape profile below is not based on this new wetland mapping because it was unavailable before release of this report. However, the data will be available in 2019 to view and download through a decision support tool web application hosted by the Nature Conservancy in Wyoming or by e-mailing Lindsey Washkoviak, Wetland Ecologist at WYNDD, [lwashkov@uwyo.edu](mailto:lwashkov@uwyo.edu).

### 3.1 Wetland Resource Description

According to mapping from the National Wetland Inventory (U.S. Fish and Wildlife Service 1984), wetland area comprises 11,636 acres, or 4% of the total land area of the LSRB study area (Table 1). This estimate excludes 1,879 acres of non-wetland features such as deep lakes, river channels, and excavated wetlands.

#### 3.1.1 Wetland Class

Palustrine Freshwater Emergent wetlands are the most common wetland class in the basin, totaling 9,526 acres and representing 82% of the wetland area (Table 1). Palustrine Freshwater emergent wetlands include irrigated hayfields, wet meadows, and emergent vegetation zones around more permanent water features such as rivers and ponds. Palustrine Forested wetlands are the second most common wetlands type and cover 1,498 acres or 12% of the wetland area. Many forested wetlands are associated with cotton wood galleries and are located along the main steam of the Little Snake River. Palustrine Scrub/Shrub wetlands account for 265 acres, and Palustrine Freshwater Pond, mainly shallow ponds, account for 316 acres. Two additional classes of Palustrine wetlands, unconsolidated bottom and unconsolidated shore, comprise the remaining 3% of wetland area.

**Table 1.** Surface area of wetlands based on NWI classifications in the LSRB.


NWI Wetland Class	Cowardin Code	Wetland Acres	% of Wetland Area
Palustrine Freshwater Emergent	PEM	9,526	81.86%
Palustrine Unconsolidated Bottom & Palustrine Unconsolidated Shore	PUB /PUS	31	0.27%
Palustrine Scrub/Shrub	PSS	265	2.28%
Palustrine Freshwater Pond	PAB	317	2.72%
Palustrine Forested	PFO	1,498	12.87%
<b>Total</b>		<b>11,636</b>	

#### 3.1.2 Water Regime

Water regime (Cowardin et al. 1979) expresses the amount of time during the year when water is present in wetlands. Seasonally and temporarily flooded wetlands are the two most common hydrologic regimes in the LSRB, representing 74% and 18% of the wetland area respectively (Figure 3). Seasonally flooded wetlands hold surface water for extended periods during the growing season but are dry by the end of the growing season in most years (Cowardin et al. 1979). Temporarily flooded wetlands hold surface water for relatively shorter periods during the



growing season. They include wetlands with hydrology dependent on alluvial groundwater and seasonal flooding along the Little Snake River and its tributaries. These water regimes represent most freshwater emergent marshes, forested wetlands, shrub wetlands and wetlands with unconsolidated bottom/shores (Table 2). Semi-permanently flooded wetlands account for 7% of the wetlands area, and all of them are as Freshwater ponds. Intermittently exposed wetlands constitute only 1% of the wetlands area (Figure 3) and are mapped as Freshwater ponds or Riverine wetlands (Table 2). Less than 1% of the wetlands are saturated (Figure 3), and they are Freshwater emergent marshes or Forested wetlands (Table 2). No wetlands within the LSRB are permanently flooded.



**Figure 3.** Surface area (acres) of wetlands classified according to NWI water regime in the LSRB.

**Table 2.** Percent of wetlands with a specific hydrologic regime in the LSRB.

Water regime		NWI Wetland Type					Palustrine Unconsolidated Bottom/Shore
		Palustrine Freshwater Emergent	Palustrine Scrub/Shrub	Freshwater Pond	Palustrine Forested	Riverine	
A	Temporarily Flooded	8%	16%	-	83%	-	36%
B	Saturated	<1%	-	-	<1%	-	-
C	Seasonally Flooded	85%	84%	-	17%	72%	64%
F	Semi-permanently Flooded	6%	-	73%	-	-	-
G	Intermittently Exposed	-	-	27%	-	28%	-

### 3.1.3 Special modifiers describing wetlands

NWI mapping includes modifier codes that identify man-made and natural alterations. Only 2% of the wetlands mapped in the LSRB have been assigned modifiers (Table 3). Modifications by beavers were identified in only 68 wetland acres. Beaver eradication has occurred throughout the basin leading to channel destabilization and down-cutting of tributaries in the basin (WBHCP 2014). At the time of study, beaver influenced wetlands occurred in drainages in the northern portion of the LSRB. Approximately 2% of wetlands (176 acres) were impounded or diked, many in intermittent drainages to retain water for livestock use. Excavated wetlands represent less than 1% (18 acres) of wetlands and were excluded from the sample frame because most of these acres were associated are gravel pits or water treatment facilities.

Irrigation was not explicitly identified as a wetland modifier in the NWI mapping codes, even though much of the land within the lower LSRB is irrigated for agricultural hay production. In addition, many modified wetlands in the LSRB were purposely created to provide waterfowl habitat or exist as a coincidence of irrigation runoff or retention. Five percent of the LSRB study area (13,374 acres) is mapped as irrigated lands (Wyoming Wildlife Consultants 2007). Although only 5% of the basin is receiving direct irrigation inputs, almost half (49%) of irrigated acres are also mapped as wetlands and over 56% of wetland acres are mapped as irrigated acres. The most common wetland type receiving irrigation inputs is freshwater emergent wetlands – 6,188 acres (65%) (Table 5). Many of these wetlands are associated with irrigation infrastructure or were created as retention ponds to provide water for livestock and wildlife.

**Table 3.** Area of wetland in acres classified according to NWI modifiers in the LSRB.

NWI Wetland Class	No Modifier		Beaver		Excavated		Impounded/diked	
	Acres	% of wetland type	Acres	% of wetland type	Acres	% of wetland type	Acres	% of wetland type
Freshwater Emergent Wetland	9506	99.9%	-	-	-	-	20	<1%
Freshwater Pond	103	31%	68	21%	13	4%	145	44%
Forested Wetland	1498	100%	-	-	-	-	-	-
Scrub/Shrub Wetland	265	100%	-	-	-	-	-	-
Unconsolidated Bottom/Shore	19	53%	-	-	5	14%	11	31%
<b>Total</b>	<b>11,391</b>	<b>98%</b>	<b>68</b>	<b>1%</b>	<b>18</b>	<b>&lt;1%</b>	<b>176</b>	<b>2%</b>




**Table 4.** Areas of irrigated wetlands in acres based on NWI classifications in the LSRB.


NWI Wetland Class	Irrigated	
	Acres	% of wetland type
Freshwater Emergent Wetland	6188	65%
Freshwater Pond	42	13%
Forested Wetland	190	13%
Scrub/Shrub Wetland	89	34%
Unconsolidated Bottom/Shore	0	0%
<b>Total</b>	<b>6,509</b>	<b>56%</b>

### 3.2 Land Ownership/Management

Land ownership in the LSRB study area is predominantly private, representing 48% of the basin, (127,033 acres) and 84% (9,849 acres) of wetland acres (Figures 4 & 5, Table 5). Lands managed by the Bureau of Land Management (BLM), State of Wyoming, and the Bureau of Reclamation comprise 38% (101,339 acres), 14% (38,079 acres), and <1% (647 acres) of the area, respectively. The Wyoming Game and Fish Commission (WGFC), in cooperation with the BLM and private landowners, manages the 37,848 acre Red-Rim Grizzly Wildlife Habitat Management Area for the co-existence of wildlife and livestock (Figure 4).



**Figure 4.** Spatial distribution of land ownership/management and wetlands within the LSRB study area.



**Figure 5.** Description of land ownership/management (acres) of the study area in the LSRB

**Table 5.** Land ownership/management of wetlands by area in the LSRB

Landowner/ Manager	Wetlands in LSRB		
	Acres	% of Basin Area	% Wetland Acres
Bureau of Land Management	1,010	0.38%	8.68%
Private	9,849	3.69%	84.64%
State	771	0.29%	6.62%
Total	11,636	4.36%	-

## 4.0 WETLAND CONDITION ASSESSMENT

### 4.1 Ecological Integrity Assessment Framework

The overarching goal of the Ecological Integrity Assessment (EIA) framework is to provide a rapid and repeatable evaluation of the ecological condition of a wetland. EIA methods were developed by NatureServe to assess the condition of wetlands across North America (Faber-Langendoen et al. 2011) and have more recently been refined by several regional wetland programs to specifically address wetland ecological condition in the Intermountain West (Rocchio 2007, Lemly and Gilligan 2012, Vance et al. 2012). We assessed condition of randomly selected wetlands in the LSRB based on EIA methods developed in Colorado by Lemly et al. (2012, 2013).

Descriptive metrics were used in the field to evaluate four attributes at each wetland: Landscape Context, Hydrologic Condition, Physicochemical Condition, and Biotic Condition. Separate disturbance indicator metrics that identify the severity of anthropogenic disturbance associated with degradation of wetland ecosystems were recorded. Metric scores for each of the four attributes were combined into an overall EIA score that can be used to describe wetlands in relation to a reference condition.

Hydrologic condition was evaluated using the Landscape Hydrology Metric (LHM) (Tibbets et al 2015) which assesses alteration to a wetland's hydrologic regime. We incorporated additional intensive assessment protocols from Colorado's EIA framework (Lemly and Gilligan 2012) including a floristic quality assessments, soil characterization, and water quality incorporated.

## **4.2 Survey Design and Evaluation of Sample Sites**

The following sections describe the survey design and process for selection of random sample sites. The steps in the survey design, were defining the target population, specifying the sample frame, choosing the sample size, and specifying the selection criteria. These methods are based on the EPA's National Aquatic Resource Survey program (Stevens & Olsen 2004; Detenbeck et al. 2005).

### ***4.2.1 Wetland Definitions for Target Population***

The target population is the set of wetlands that we want to characterize in the LSRB. Our wetland target population consisted of the six classes of Palustrine wetlands that we used in the landscape profile (see Table 1). Palustrine wetlands can be situated shoreward of lakes or river channels, on floodplains, in locations isolated from water bodies, in depressions, or on slopes. The target population included all palustrine wetlands within the LSRB study area and excluded non-wetland features such as deepwater lakes (Lacustrine system) and stream channel bottoms (Riverine system) (Table 6). We also set a minimum size threshold of at 0.1 hectare and a minimum width of 10 m.

### ***4.2.2 Sample Frame and Classification***

The sample frame is a digital representation of the target population. The digital NWI polygon dataset (U.S. Fish and Wildlife Service 1984) is a complete representation of the target population, but it contains a degree of detail that makes it very difficult to use without grouping NWI codes into wetland subgroups. We simplified the sample frame by grouping the NWI Cowardin et al (1979) codes into three target groups: 1) Wet meadow; 2) Emergent Marsh; and 3) Riverine Shrubland. We the crosswalk each target wetland subgroup to the Ecological Systems classification (Comer et al. 2003, Appendix A). Classification by Ecological Systems is the dominant system used regionally for identifying wetland types in the field and provides a valuable system for defining landscape units by biotic (e.g., plant community) and abiotic (e.g., geologic, hydrologic, elevation) criteria (Lemly and Gillian 2012, Newlon et al. 2013). NWI codes were also crosswalk to Hydrogeomorphic (HGM) Classification (Brinson 1993, Adamus 2004) (Table 6).

**Table 6.** Target wetland subgroups classified by Cowardin, Hydrogeomorphic (HGM), and Ecological Systems used in the LSRB.

Targeted Wetland Subgroups	NWI Codes	HGM	NWI Codes	Ecological System
Wet Meadows	PEMB, PEMA, PEMC	Slope, Depression	PEMB, PEMA, PEMC	Rocky Mountain Alpine-Montane Wet Meadow, Irrigated Wet Meadow (not an official Ecological System)
Emergent Marsh	PEMFh, PUSCh, PUSC, PABG, PUSAh, PEMCh, PUSC, PABGh, PABF, PABGh, PEMF, PEMAh	Depression, Riverine	PEMFh, PUSCh, PUSC, PABG, PUSAh, PEMCh, PUSC, PABGh, PABF, PABGh, PEMF, PEMAh	Western North American Emergent Marsh
Riverine Shrubland	PABFb, PSSB, PABGb, PSSA, PSSC, PFOA	Riverine	PABFb, PSSB, PABGb, PSSA, PSSC, PFOA	Rocky Mountain Lower Montane-Foothill Riparian Woodland and Shrubland

#### 4.2.3 Sample Size and Selection Criteria for Site Evaluation

The target sample size was 75 sites selected from the sample frame, divide across the three target wetland subgroups. Sample sites were randomly selected from the sample frame using a Generalized Random Tessellation Stratified (GRTS) survey design for a finite resource (Stevens et al. 2004, Stevens and Jensen 2007). GRTS sampling was performed using R package spsurvey (Olsen and Kincaid 2009, R Development Core Team 2014).

After potential sample sites were randomly selected, and prior to field sampling, a desktop site evaluation was performed to determine: 1) whether a wetland was likely present, based on examination of aerial imagery (USDA Farm Service Agency 2009); and 2) land ownership/management status (private, state, federal). Permission was then sought to access sample sites.

Potential sample sites that met one of the following conditions were withdrawn from potential sample sites *before* field sampling:

1. Wetland type: the wetland at the site appeared to not belong to the target wetland group that the site was chosen to represent.
2. Size: the wetland area did not meet the minimum 0.1-hectare area threshold or 10-meter width threshold required for sampling

3. Minimum distance: the wetland was within 500 meters of another sample location of the same target subgroup.
4. Permission denied: permission to access the site was denied by the landowner.

Sites that remained after the initial review were visited and were evaluated by field crews before assessment. Sites that met one of the following conditions were withdrawn from the sample frame before field assessment:

5. Wetland type: the wetland did not belong to the target wetland group that the site was chosen to represent. The field crew used a key to ecological systems (Appendix A) for this evaluation
6. Access issues: permission was granted by landowner, but the point could not be safely accessed at the time of sampling. Sites were rejected if they were more than 2 miles from the vehicle for efficiency and safety of the field crew.
7. Depth: the wetland exceeded the maximum depth threshold of 1 meter and the assessment area could not be repositioned to a location that met our size criterion.
8. Hayed before sampling: all of the vegetation was cropped from the site prior to sampling, so that plant identification was not possible.
9. Not a wetland: the sample location did not meet our operational definition of a wetland (Appendix B) or no wetland was present due to mapping error

If a site was withdrawn from the set of potential sample sites, it was replaced with the next site from the sequential list generated by the GRTS site selection.

In addition to the random survey sites, we identified 3 to 4 reference wetlands from each target wetland subgroup that represented “least disturbed” condition based on professional judgment of regional wildlife managers or the field crew. We used the definition provided by Stoddard et al. (2006) for least disturbed condition: “in the best available physical, chemical and biological habitat conditions given today’s state of the landscape”.

### **4.3 Field Methods**

Field methods were based on EIA protocols developed by Lemly et al. (2013). In addition, we collected data on soils and vegetation to supplement the EIA protocol. These assessments required a half a day or less to complete at each site. Detailed field data forms are included in Appendix C and our field manual is available upon request.

#### **4.3.1 Wetland Assessment Area (AA)**

The field crew applied the EPA’s National Wetland Condition Assessment (U.S. Environmental Protection Agency 2016) methodology for establishing an assessment area (AA) at each wetland site. When possible a standard 40 m radius circular AA was established. If the site configuration did not accommodate a circular AA of this size, the crew adjusted the AA to a rectangular or irregular shape of at least 1000 m<sup>2</sup> (0.1 ha) and 10 m wide. The AA boundary was

marked with flagging to aid with data collection. A 500-m buffer was established from the perimeter of each AA. Standard descriptions of each wetland included: UTM coordinates, wetland classification, presence or signs of wildlife, and photos of the buffer and AA.

#### 4.3.2 Ecological Integrity Assessment of Wetland Sites

After the AA was established, each wetland was assessed based on the EIA manual and field forms (see Appendix C) adapted from Lemly et al. (2013). The principal attributes and metrics that were measured in this study are summarized in Table 7.

**Table 7.** EIA attributes and metrics used for wetland assessments in the LSRB.

Attributes	Indicators and Metrics
Landscape Context	<ul style="list-style-type: none"> <li>• Landscape Fragmentation</li> <li>• Buffer Extent</li> <li>• Buffer Width</li> <li>• Buffer Condition</li> </ul>
Hydrologic Condition*	<ul style="list-style-type: none"> <li>• Water Source</li> <li>• Hydrologic Connectivity</li> <li>• Alteration of Hydroperiod</li> </ul>
Physicochemical Condition	<ul style="list-style-type: none"> <li>• Water Quality</li> <li>• Algal Growth</li> <li>• Substrate/soil Disturbance</li> </ul>
Biological Condition	<ul style="list-style-type: none"> <li>• Relative Cover of Native Plant Species</li> <li>• Absolute Cover of Noxious Weeds</li> <li>• Absolute Cover of Aggressive Native Species</li> <li>• Mean C</li> <li>• Structural Complexity</li> </ul>

\*Field data were collected for the EIA hydrology metrics using the Colorado EIA method, however, we used Landscape Hydrologic Metric in place of the Colorado EIA method for scoring wetland condition.

#### 4.3.3 Landscape Hydrology Metric (LHM)

Hydrology is broadly characterized as the movement, distribution, timing, and quality of water across the landscape. Hydrology is the primary driver of the processes that establish and maintain wetlands, including ecological, physical, and chemical processes that sustain ecosystem functions and associated services and values to people (Mitsch and Gossilink 2000). Therefore, it is important to identify alterations to the natural hydrologic regime that may detrimentally affect the structure and function of a wetland. Identifying alterations to natural wetland hydrology can be a challenge because significant alterations such as major dams or ditches may

not be evident during a single site visit or are located outside the 500m buffer surrounding the AA. In addition, it can be difficult to identify a wetland’s water source when the wetland is supported or created by hydrologic alterations, such as leaky dams or canals.

We used the Landscape Hydrology Metric (LHM) (Tibbets et al. 2015), instead of the hydrology component of the Colorado EIA method (Lemly et al. 2013), to calculate the hydrologic condition metrics. LHM incorporates landscape-level data identifying alterations to hydroperiod and water source, along with field data characterizing wetland soils. LHM relies on descriptive criteria from submetrics to assign a rank from 5 to 0 (Table 8). Historic wetlands (score = 5) were defined in this study as wetlands without evidence of hydrologic alteration, whereas created wetlands (score = 0) are dependent on hydrologic alteration.

**Table 8.** Landscape Hydrology Metric scoring criteria.

Hydrologic Category	LHM Score	Landscape Hydrology Metric Criteria
Historical Wetland	5	No alterations to hydrology identified, natural water source or no observed natural water source but histic soil layer present.
Hybrid Wetland in landscape with site-level hydrologic alterations	4	Site-level hydrologic alteration, natural water source identified or no observed natural water source but histic soil layer present.
Hybrid Wetland in landscape with basin-wide hydrologic alterations	3	Basin-wide hydrologic alteration (major dam present) and direct hydrologic connectivity to natural water source observed. No histic soil layer observed.
Supported Wetland with natural water source	2	Basin-wide hydrologic alteration (major dam present), landscape position is in depression with natural water source potential, however, dominant water source is unclear due to presence of large canals. No histic soil layer observed.
Supported Wetland- Irrigation Dependent Depression	1	Hydrologic alteration identified, landscape position is in depression. Irrigation is likely dominant water source. No histic soil layer observed.
Created Wetland - Irrigation Dependent	0	Hydrologic alteration identified, no natural water source identified. Irrigation is exclusive water source. No histic soil layer observed.

#### 4.3.4 Vegetation Assessments

We used a plotless sample design to collect vegetation data using methods described in Lemly et al. (2012). Species searches were limited to no more than 1 hour at each site. Vascular plant species were identified using Dorn (2001) and regional keys including Johnston (2001), Skinner



(2010), and Culver and Lemly (2013). Species names were taken from the WYNDD database. Unknown plant specimens were pressed in the field and saved for later identification. The percent cover of each species, including that of unidentified specimens, was estimated over the entire AA.

#### ***4.3.5 Soils***

We dug 1 to 2 soil pits within each AA. The first soil pit was placed in a representative location close to the AA center excluding those areas covered completely by water. An additional pit was dug if there was a high degree of variability within the site. We recorded GPS waypoints at each soil pit and then marked the location on a map. Pits were dug to a depth of 40 cm (about one shovel length) when possible. The core was removed and laid next to the pit, ensuring all horizons were intact and in order. We recorded the following information from each horizon: 1) color (based on a Munsell Soil Color Chart (2013)) of the matrix and any redoximorphic concentrations (mottles and oxidized root channels) and depletions; 2) soil texture; and 3) any other specifics about the concentration of roots, the presence of gravel or cobble, or other unusual soil features. Hydric soil indicators were identified based on guidance from the Interim Regional Supplement to the U.S. Corps of Engineers Wetland Delineation Manual: Western Mountains, Valleys, and Coast Region (2008) and the National Resources Conservation Service (NRCS) Field Indicators of Hydric Soils in the United States (Vasilas et al. 2010).

#### ***4.3.6 Surface Water Characterization***

We estimated percent cover and interspersion (patch complexity) of open water within the AA. The water depth range and average were recorded within the AA.

### **4.4 Data Analysis**

#### ***4.4.1 Data Management***

All field data were entered into relational databases that were developed using Microsoft Access and/or ArcGIS 10.3 platforms. Data were then proofed to correct any errors prior to analysis. The data are stored at the Wyoming Natural Diversity Database.

#### ***4.4.2 Ecological Integrity Assessment***

To be effective tools, ecological assessment metrics should provide information about the integrity of major ecological attributes in relation to a gradient of disturbance or stressors. We evaluated performance of each EIA metric based on methods used to refine stream and wetland condition indices (Stoddard et al. 2008, Deller et al. 2010, Faber-Langendoen et al. 2011). Evaluation of EIA methods and scoring was a vital step to ensure the EIA methods we selected were relevant and effective for assessing wetland condition in Wyoming. The applicable range of each metric was determined by examining histograms depicting ranges and distributions of scores. We evaluated metric redundancy by calculating Spearman's rank correlation coefficients among all metrics. None of the metrics within an attribute category were found to be highly correlated (as determined by a coefficient value of  $r > 0.8$ ).



We calculated EIA scores and thresholds based on EIA methods used in Colorado (Lemly and Gillian 2012). Refer to Appendix D for a detailed description of scoring formulas and thresholds with ranks ranging from A-D. Ideally, wetlands that are ranked “A” are those in minimally disturbed condition (MDC), representing the best approximation of “naturalness” or a high degree of “biological integrity” on the landscape (Stoddard et al. 2006). However, reference wetland condition in the LSRB was defined as least disturbed condition (LDC), meaning “in the best available physical, chemical and biological habitat conditions given today’s state of the landscape” (Stoddard et al. 2006). Because LDC can differ from MDC, the biological integrity of our A-ranked sites may not reflect the sites’ fullest potential for biological integrity.

We created a cumulative distribution function (CDF) plot to display EIA scores estimated for wetlands across the entire sample frame in the LSRB. CDF plots use scores from the random sample to create a probability plot for the entire basin. CDF plots are useful to estimate the cumulative proportion of the resource (wetlands) estimated to have at least a certain EIA score (Whittier et al. 2002). EIA rank thresholds were superimposed on the CDF plot to facilitate interpretation of the cumulative number of wetlands within each rank. Cumulative distribution functions were calculated using R software package version 3.3.3 (R Development Core Team 2014) available from the *spsurvey* library.

#### ***4.4.3 Floristic Quality Assessment (FQA)***

Floristic Quality Assessment (FQA) uses plant community composition as an indicator of ecological condition. The FQA method assesses the degree of human caused disturbance based on the proportion of “conservative” plants present. “Coefficients of conservatism” (C-values) are the foundation of FQA. C values range from 0 to 10 and represent an estimated probability that a plant is likely to occur in a landscape relatively unaltered from conditions that existed before European settlement (Swink and Wilhelm 1979, 1994). A C-value of 10 is assigned to plant species having low tolerance for habitat degradation and are restricted to relatively unaltered areas, whereas a 0 is assigned to plant species with a wide tolerance to human disturbance (Rocchio 2007). Species with low C values may be found in relatively unaltered areas, but they also grow in altered areas. Once C-values have been assigned for a given region or area, they can then be used to calculate a number of FQA indices such as the average C-value of a site (Mean C) and the Floristic Quality Assessment Index (FQAI) (Swink and Wilhelm 1979, 1994). C-values were developed for Wyoming in 2017 and have been incorporated into data analysis (Washkoviak et al. 2017)

We calculated Mean C, total species richness, and the numbers of native and non-native species from the species lists compiled at each wetland site. Mean C for the site calculated by summing the C-values of the plant species found at each site, and then dividing by the number of species.

## 5.0 WETLAND CONDITION ASSESSMENT RESULTS

### 5.1 Characteristics of Sampled Wetlands

Seventy wetlands, including 3 reference sites, were sampled in 2016. We evaluated 259 sites of the three target subgroups (freshwater emergent marsh, riverine shrublands, and wet meadows) using the defined selection criteria, but not all sites were sampled or included in the study (Table 9). One site was rejected from evaluation because it was located outside of our sample area boundary due to a mapping error. Permission was denied at 115 sites and six sites did not meet minimum the distance threshold (Table 10). Access issues due to limited roads and rough terrain accounted for 21 site rejections. Fifty sites were rejected in the field before sampling because they did not meet our operational definition of a wetland.


**Table 9.** Number of wetland sites in each target wetland subgroup evaluated, sampled, and rejected during the study by wetland subgroup.

Target wetland subgroup	# Sites in survey design	# Sites evaluated	# Random survey sites sampled	# Sites rejected (see table 10)
Freshwater Emergent Marsh	100	86	18	68
Riverine Shrubland	89	80	24	56
Wet meadows	110	93	25	68
<b>Total</b>	<b>299</b>	<b>259</b>	<b>67</b>	<b>192</b>

**Table 10.** Reasons for rejection of wetlands in the LSRB during the study.

Target wetland subgroup	Rejection Cause			
	Permission denied	Minimum distance	Access Issue	Not a wetland
Freshwater Emergent Marsh	39	3	7	19
Riverine Shrubland	37	3	5	11
Wet meadows	39	0	9	20
<b>Total</b>	<b>115</b>	<b>6</b>	<b>21</b>	<b>50</b>

We obtained permission to sample 31% of the sites selected in the random survey design. Thirty-nine (56%) sampled wetlands were on private lands, 15 (23%) on State lands, and 16 (21%) on lands administered by the BLM (Figure 6). The percentage of the sampled points on private lands (56%) was less than the percentage of the potential, randomly selected points on private lands (70%), revealing a bias in sampling toward public-land sites.



**Figure 6.** Percentage of points selected in the original survey design versus sites sampled by landowner/manager

### 5.1.1 Characteristics of Sampled Wetland Subgroups

A field key developed for wetlands in Wyoming was used to classify each sampled wetland according to an ecological system (Appendix A). After completion of the field survey, we summarized the general characteristics of each of the three target wetland subgroups. The descriptions below include specific observations made during field sampling in the LSRB combined with more general information from the ecological system key.

#### Riverine Shrubland

Riverine shrublands are typically distributed along the Little Snake River and its tributaries within the LSRB. Riverine shrublands are dominated by a shrub overstory of *Salix* sp., *Ribes* sp. and *Pentaphylloides floribunda* with a mesic to hydric meadow understory vegetation of native and non-native grasses and forbs such as *Poa pratensis*, *Phleum pretense*, *Carex utriculata*, *Juncus balticus*, *Mentha arvensis*, and *Cirsium arvense*. Many are associated with historical floodplains and receive water from overbank flooding and alluvial aquifers. Some Riverine shrubland complexes higher in the basin are associated with peat soil layers, likely relics of historic beaver activity in the basin (Knight et al. 2014).



**Figure 7.** Riverine shrubland wetlands in the LSRB.

### **Freshwater Emergent Marsh**

Freshwater marshes and ponds include riverine oxbows, created ponds receiving irrigation inputs, and some areas along the shorelines of major reservoirs within the basin. Marshes characteristically have central areas that are frequently flooded and surrounded by increasingly drier zones. The central area is dominated by hydrophytic species such as *Eleocharis palustris*, *Polygonum amphibium*, and *Hippuris vulgaris*. Dominant species in the surrounding dryer zones include *Hordeum jubatum*, *Phalaris arundinacea*, *Juncus balticus*, *Alopecurus arundinaceus*, and *Cirsium arvense*.



**Figure 8.** Freshwater emergent marsh wetlands in the LSRB.

### **Wet meadows**

Wet meadows are wetlands dominated by native and non-native herbaceous vegetation, often within floodplains with a high-water table and/or artificial overland flow (irrigation). These sites typically lack prolonged standing water. Graminoids typically comprise the greatest canopy



cover. Common native species in the LSRB include *Juncus balticus*, *Carex nebrascensis*, *Achillea millefolium*, and *Deschampsia cespitosa*. Non-native hay grasses such as *Poa* spp., *Alopecurus* sp, *Phleum pretense*, and *Agrostis stolonifera* are often abundant within wet meadows. Standing water less than 0.1 ha in size can exist within wet meadows and may sustain emergent marsh vegetation but it is not the dominant Ecological System.



**Figure 9.** Wet meadow wetlands in the LSRB.

## 5.2 Characterization of Wetland Vegetation

### 5.2.1 Species Diversity of Wetland Vegetation

Plant surveys identified 273 taxa of vascular plants at the 70 wetlands sampled. Thirty-seven were identified only to genus and two more only to family. Two species were unidentifiable because diagnostic floristic parts required for species identification were absent at the time of sampling. The remaining 232 taxa were identified to the species and subspecies level and represent 8% of Wyoming's flora (Dorn 2001).

The two most common plant species found at wetlands sampled in the LSRB were native. Nebraska Sedge (*Carex nebrascensis*) and Baltic Rush (*Juncus balticus*) were found at 76% and 73% of sites respectively (Table 11 & 12). Many non-native species are commonly planted lawn and pasture grasses such as Kentucky Blue Grass (*Poa pratensis*), Common Timothy (*Phleum pratense*), Spreading Bent (*Agrostis stolonifera*), Creeping Meadow Foxtail (*Alopecurus arundinaceus*), Dandelion (*Taraxacum officinale*), and White Clover (*Trifolium repens*). Canada thistle (*Cirsium arvense*), a state designated noxious weed, was found at 36% of sites.

**Table 11.** Ten most common plant species documented at sampled wetland in the LSRB.

Species	% of Sites	Wetland Status	Nativity	WY C Value	Common Name
<i>Carex nebrascensis</i>	76%	OBL	Native	4	Nebraska Sedge
<i>Juncus balticus</i>	73%	FACW	Native	3	Baltic Rush
<i>Poa pratensis</i>	71%	FAC	Exotic	0	Kentucky Blue Grass
<i>Taraxacum officinale</i>	69%	FACU	Exotic	0	Common Dandelion
<i>Trifolium repens</i>	53%	FACU	Exotic	0	White Clover
<i>Phleum pratense</i>	53%	FACU	Exotic	0	Common Timothy
<i>Cirsium arvense</i>	50%	FACU	Exotic	0	Canadian Thistle
<i>Achillea millefolium</i>	49%	FACU	Native	4	Common Yarrow
<i>Mentha arvensis</i>	49%	FACW	Native	4	American Wild Mint
<i>Deschampsia cespitosa</i>	49%	FACW	Native	5	Tufted Hair Grass
<i>Potentilla anserina</i>	47%	OBL	Native	4	Common Silverweed

**Table 12.** Frequencies of native and non-native species encountered at the sites sampled in the LSRB.

Native		Non-Native	
Species	% of Sites	Species	% of Sites
<i>Carex nebrascensis</i>	76%	<i>Poa pratensis</i>	71%
<i>Juncus balticus</i>	73%	<i>Taraxacum officinale</i>	69%
<i>Deschampsia cespitosa</i>	49%	<i>Trifolium repens</i>	53%
<i>Mentha arvensis</i>	49%	<i>Phleum pratense</i>	53%
<i>Achillea millefolium</i>	49%	<i>Cirsium arvense</i>	36%
<i>Potentilla anserina</i>	47%	<i>Agrostis stolonifera</i>	47%
<i>Equisetum laevigatum</i>	46%	<i>Alopecurus arundinaceus</i>	40%
<i>Carex utriculata</i>	43%	<i>Plantago major</i>	16%
<i>Cirsium scariosum</i>	40%	<i>Bromus inermis</i>	16%
<i>Epilobium ciliatum</i>	39%	<i>Lactuca serriola</i>	10%

### 5.2.2 Floristic Quality Assessment

Across all wetland types, non-native species comprised 15-32% of the mean relative cover at sampled wetlands in the LSRB (Table 13). Mean C values across wetland sites was 3.44 and ranged between (1.67 -7). Mean C-values for native species fell between 4 and 7, indicating most species observed had some degree of specificity for unaltered habitat but had a moderate tolerance to disturbance. Riverine Shrublands had the highest Mean C, mean species richness, and FQI, even though non-native species averaged 20% of the relative cover at these sites.

**Table 13.** Floristic quality assessment indices calculated for wetlands in the LSRB.

FQA Indices	Riverine Shrubland		Freshwater Emergent Marsh		Wet Meadow		Overall	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
<b>Total species richness</b>	36.32	10.03	22.36	11.69	25.48	8.78	27.44	11.13
<b>Native species richness</b>	25.57	9.12	15.23	7.08	17.59	7.23	19.01	8.69
<b>Non-native species richness</b>	6.89	3.83	4.85	3.08	5.21	2.18	5.57	3.05
<b>Mean C of all species</b>	3.59	.81	3.43	.94	3.34	.7	3.44	.80
<b>Mean C of native species</b>	4.60	.42	4.28	.68	4.48	.43	4.45	.53
<b>FQI of all species</b>	21.71	6.83	15.09	4.05	16.91	5.28	17.64	5.94
<b>FQI of native species</b>	23.15	5.96	15.91	4.29	18.42	4.86	18.91	5.70
<b>Adjusted FQI</b>	.30	.09	.28	.07	.28	.08	.29	.08
<b>Relative % cover native species</b>	79.47	.09	84.88	.14	67.51	.18	76.22	.16
<b>Absolute % cover Non-native species</b>	26.52	22.11	8.75	8.55	33.87	19.18	24.42	20.47
<b>Absolute % cover noxious species</b>	3.63	5.00	2.69	5.00	1.31	1.39	2.63	4.23

### 5.2.3 Plant Species of Conservation Concern

The Wyoming Natural Diversity Database (WYNDD) develops and maintains lists of species in Wyoming that are rare, endemic, disjunct, threatened, or otherwise biologically sensitive (WYNDD 2017). The Wyoming Plant Species of Concern List (SOC) and the Wyoming Plant Species of Potential Concern List (SOPC) show the vascular plant species, subspecies and varieties that meet one of these one or more of these criteria.

Three state-critically imperiled (S1), 15 state-imperiled (S2) species, six state-imperiled/vulnerable (S2S3) species, and 41 state-vulnerable (S3) species were found at the sampled sites in the LSRB (Table 14). No globally vulnerable (G ranked) species were found during sampling.

**Table 14.** Plant species of concern identified at sampled wetlands within the LSRB.

<b>S1 Species</b>	<b># Sites</b>
<i>Najas guadalupensis</i>	1
<i>Packeria pseud aurea</i> var. <i>flavula</i>	2
<i>Spirodela polyrhiza</i>	4

<b>S2 Species</b>	<b># Sites</b>
<i>Carex retrorsa</i>	1
<i>Cerastium brachypodium</i>	7
<i>Ceratophyllum demersum</i>	2
<i>Danthonia californica</i>	3
<i>Elodea canadensis</i>	2
<i>Epilobium palustre</i> var. <i>palustre</i>	1
<i>Hypericum formosum</i>	5
<i>Juncus alpinoarticulatus</i>	1
<i>Ranunculus gmelinii</i>	4
<i>Rumex densiflorus</i>	2
<i>Sidalcea candida</i>	9
<i>Sidalcea neomexicana</i>	25
<i>Sium suave</i>	3
<i>Stuckenia vaginata</i>	4
<i>Thermopsis montana</i>	4
<i>Typha angustifolia</i>	3

<b>S2/3 Species</b>	<b># Sites</b>
<i>Conioselinum scopulorum</i>	1
<i>Glyceria elata</i>	11
<i>Polemonium occidentale</i>	7
<i>Ranunculus acriformis</i>	1
<i>Thalictrum dasycarpum</i>	1
<i>Utricularia macrorhiza</i>	1

<b>S3 Species</b>	<b># Sites</b>
<i>Aconitum columbianum</i>	1
<i>Alisma triviale</i>	5
<i>Angelica pinnata</i>	5
<i>Astragalus bodinii</i>	1
<i>Calamagrostis stricta</i>	1
<i>Carex athrostachya</i>	1
<i>Carex canescens</i>	1
<i>Carex simulata</i>	2

<b>S3 Species - continued</b>	<b># Sites</b>
<i>Catabrosa aquatica</i>	8
<i>Cirsium scariosum</i>	28
<i>Dracocephalum parviflorum</i>	1
<i>Epilobium lactiflorum</i>	1
<i>Galium trifidum</i>	10
<i>Glaux maritima</i>	5
<i>Glyceria grandis</i>	9
<i>Hackelia patens</i>	3
<i>Hippuris vulgaris</i>	7
<i>Juncus ensifolius</i> var. <i>ensifolius</i>	21
<i>Juncus longistylis</i>	2
<i>Limosella aquatica</i>	2
<i>Myriophyllum sibiricum</i>	5
<i>Parnassia palustris</i> var. <i>montanensis</i>	1
<i>Plantago eriopoda</i>	6
<i>Polygonum ramosissimum</i>	2
<i>Potamogeton gramineus</i>	1
<i>Potentilla anserina</i>	33
<i>Potentilla biennis</i>	1
<i>Prunella vulgaris</i>	10
<i>Ranunculus abortivus</i>	1
<i>Ranunculus sceleratus</i> var. <i>multifidus</i>	2
<i>Ribes inerme</i>	2
<i>Rorippa palustris</i>	3
<i>Rudbeckia laciniata</i> var. <i>ampla</i>	1
<i>Rumex aquaticus</i> var. <i>fenestratus</i>	3
<i>Rumex maritimus</i>	1
<i>Schoenoplectus acutus</i>	13
<i>Scirpus microcarpus</i>	15
<i>Scutellaria galericulata</i>	1
<i>Sisyrinchium montanum</i>	2
<i>Solidago gigantea</i>	5
<i>Sparganium emersum</i>	3
<i>Stachys palustris</i> var. <i>pilosa</i>	5
<i>Stellaria longifolia</i>	9
<i>Stuckenia pectinata</i>	5
<i>Suaeda calceoliformis</i>	2
<i>Triglochin palustris</i>	3



### 5.3 Wildlife Observations

Wildlife observations were recorded opportunistically during wetland sampling. Wildlife or wildlife sign were observed at 96% of sites, including 42 unique observations of birds, nine amphibians, seven native ungulates, and five mammals (Table 15). Formal bird surveys by the USFWS and WGFD are ongoing in the study area.

**Table 15.** Wildlife observations made during wetland sampling effort in the LSRB.

	Wildlife	# of Sites		
		Visual	Tracks or Vocalization	Nest or Den
<b>Ungulates</b>	Mule deer	1	2	
	Elk	1	3	
<b>Birds</b>	Redwing blackbird	13		
	Sand hill crane	2		
	Ducks	7		
	Killdeer	4		
	American coot	1		
	Gull	1		
	Wilson's Snipe	4		
	Swallow	4		
	Marsh wren	1		
	Northern harrier	1		
	Red tailed hawk	1		
	White pelican	1		
<b>Mammals</b>	Beaver	1	1	1
	Muskrat			1
	Striped skunk	1		
<b>Amphibians</b>	Tiger salamander	1		
	Unknown frog species	7	1	
<b>Reptiles</b>	Garter Snake	2		
<b>Crustacean</b>	Crawfish	2		


## 5.4 Wetland Condition Assessment

### 5.4.1 Ecological Integrity Assessment of Sampled Wetlands

Ecological integrity assessment (EIA) scores from the 70 sampled wetlands ranged from 1.97 to 4.63 out of a possible range of 1.0 - 5.0. We established four wetland condition categories based on values defined in Appendix D:

- A (4.5 - 5.0) = At or near reference condition (no or little human impact-see Section 4.4.3)
- B (3.5 - < 4.5) = Level of disturbance indicates slight departure from reference condition
- C (2.5 - < 3.5) = Level of disturbance indicates moderate deviation from reference condition
- D (< 2.5) = Level of disturbance indicates severe deviation from reference condition

Seven percent of the 70 study sites in the LSRB were ranked “A,” 60% were ranked “B,” 24% were ranked “C,” and 9% were ranked “D” (Fig. 5). All three target wetland subgroups were dominated by B-ranked wetlands. Riverine Shrublands had the highest proportion of A-ranked sites and no D-ranked sites, indicating overall lower disturbance relative to other wetland types. Wet meadows and emergent marshes scored a C or below for 30% and 50% of sites, respectively.



**Figure 10.** Sixty-seven percent of the wetlands in the LSRB received an EIA score of B or above. EIA scores were lowest for emergent marshes, indicating 50% of these wetlands surveyed received a C or below, indicating moderate to significant departure from reference

EIA scores were derived from 4 attributes: landscape context, biotic condition, physicochemical condition, and the Landscape Hydrology Metric. Sixty percent of wetlands sampled received a landscape context rank of A (Table 16) meaning that over half of wetlands sampled existed in landscapes where there was little landscape fragmentation within 500 meters of the sample site. No wetlands had a landscape context rank of D. No wetlands received a biological condition ranking of A, and, 46% and 47% of sampled sites received a ranking of B or C, respectively. Riverine Shrublands biotic ranks were generally higher compared to other wetland subgroups. The Physicochemical condition rankings were generally A or B, with the exception of 7 C-ranked wetlands (mainly emergent marshes). No wetlands had Physicochemical Ranks of D. Frequencies of LHM classifications within wetland subgroupings are shown at the bottom of Table 16 for comparison to the other EIA attribute ranking frequencies.

**Table 16.** Ranks for each EIA attribute by wetland subgroup for the LSRB.

Wetland Subgroup	EIA Landscape context rank			
	A	B	C	D
Emergent Marsh	11	8	3	0
Riverine Shrubland	15	4	0	0
Wet Meadow	16	9	4	0
<b>Total</b>	<b>42</b>	<b>21</b>	<b>7</b>	<b>0</b>

Wetland Subgroup	EIA Biotic condition rank			
	A	B	C	D
Emergent Marsh	0	10	12	0
Riverine Shrubland	0	14	3	2
Wet Meadow	0	8	18	3
<b>Total</b>	<b>0</b>	<b>32</b>	<b>33</b>	<b>5</b>

Wetland Subgroup	EIA Physicochemical condition rank			
	A	B	C	D
Emergent Marsh	7	11	4	0
Riverine Shrubland	14	4	1	0
Wet Meadow	21	6	2	0
<b>Total</b>	<b>42</b>	<b>21</b>	<b>7</b>	<b>0</b>


Wetland Subgroup	LHM Hydrology classification			
	Historical	Hybrid	Supported	Created
Emergent Marsh	3	11	6	2
Riverine Shrubland	15	3	0	1
Wet Meadow	19	4	1	5
<b>Total</b>	<b>37</b>	<b>18</b>	<b>7</b>	<b>8</b>

### 5.4.2 Estimate of Wetland Condition for targeted wetlands in the LSRB

We used EIA scores and condition category thresholds to estimate the ecological condition of wetlands throughout the LSRB study area. Cumulative distribution function (CDF) estimates were derived from the sample design to estimate the number of wetlands each sample site represented across the total sample frame in the basin. Percent and standard error of number of wetlands within each ranking category were calculated and are shown in Table 17. The CDF plot is not linear, indicating that estimated EIA scores are not evenly distributed across the wetland population (Figure 11). Confidence intervals vary along the plot and are widest at the lowest scores. Based on CDF analysis, 7% of wetlands in the LSRB are estimated as A-ranked, 65% B-ranked, 17% C-ranked and 11% D-ranked wetlands (Table 17). An assumption of the CDF analysis is that data were obtained from a random sample representative of the wetland population in the study area. However, the sample size was limited because of either landowner denial of access (38% of wetlands) or other rejection criteria (26%).

**Table 17.** Population estimate of EIA ranks for wetlands in the LSRB. Observed = percent of sampled sites within each rank; Estimate = percent of wetland number extrapolated using 67 wetlands from the sample frame

EIA Rank	Observed in Sample	Estimated for Basin	95% Confidence Interval
A	7%	7%	0-14%
B	60%	65%	55-76%
C	24%	17%	10-24%
D	9%	11%	4-18




**Figure 11.** Cumulative distribution function of wetland EIA scores with 95% CI shown. Graph is the cumulative proportion of wetlands (y-axis) with EIA scores at or below values on the x axis.

### 5.4.3 Landscape Hydrology Metric

The Landscape Hydrology Metric (LHM) is an assessment of alteration to hydrologic regime of sampled wetlands. LHM incorporates landscape-level data identifying alterations to hydroperiod and water source, along with field data characterizing wetland soils. LHM categories (historical, hybrid, supported, created) are defined in Table 8 above

Fifty-three percent of the wetlands sampled in the LSRB were historical and 26% were considered hybrid, while 10% were considered supported and 11% created. (Figure 12). Wet meadows had both the highest percentage of historical sites (65%) as well as the highest percentage of sites created by hydrological alterations (17%) (Table 16).




**Figure 12.** Proportion of total wetland sites based on categories for the Landscape Hydrology Metric.

### 5.4.4 Indicators of disturbance

Potential indicators of disturbance include natural phenomena or human caused land management impacts that have the ability to stress a wetland or reduce its ecological condition. These indicators can be used to identify the most prevalent impacts affecting wetland health in a given area and can help land managers change and address disturbances that are under their control. We recorded indicators of disturbance within a 500-meter wide buffer around the wetland and within the wetland boundary. These indicators were later grouped into categories based on disturbance type. A full list is available in Appendix E.

The most common potential indicators of disturbance the LSRB are listed in Figure 13. Almost all (97%) of sites sampled had non-native species present. Roads were recorded within 500 meters of 60 (86%) sites. Roads lead to fragmentation and can change the lateral movement of water by both blocking and diverting flow. Lightly used 2-tracks or farm roads and gravel roads were the


most common type of road. Soil degradation from livestock and wild horses occurred at 27 (69%) sites. Light to moderate grazing by livestock or native ungulates was present at 41 (57%) sites. Only 5 wetlands (< 1%) were being impacted by heavy grazing (plants grazed less than 3 inches). Irrigation infrastructure, including ditches, berms, and head gates that change the flow of water, was the third most common stressor impacting 33 (47%) sites. In Wyoming irrigation infrastructure can both create and degrade wetlands.



**Figure 13.** Potential indicators of disturbance observed across all wetlands in the Little Snake River Basin.

## 6.0 LITTLE SNAKE FOCAL AREAS

After sampling and reviewing condition assessment results, we found that the LSRB study area can be divided into three distinct areas that have different landowner/management patterns, land uses, wetland types, and potential patterns of disturbance. We are identifying these areas in this report as the Upper Basin, the Floodplain, and the Muddy Creek Wetlands Project focal areas (Figure 14). In this section, we highlight key differences between focal areas that can be used to complement the interpretation of the basin-scale results.



**Figure 14.** Focal areas within the LSRB showing landownership/management

### 6.1 The Muddy Creek Wetlands Project

The Muddy Creek Wetlands Project (MCWP) is located near Dad, WY along Muddy Creek 20 miles north of the town of Baggs, WY. The MCWP is the largest constructed wetland complex in Wyoming, consisting of 32 different ponds creating approximately 1500 acres of wetland and riparian habitat along 7 miles of Muddy Creek. The MCWP was constructed to provide habitat for migrating and breeding birds (Wyoming Game and Fish Department 2018).

Wetlands in the MCWP were included in the survey design but no random sample sites were selected there. We targeted two wetlands, an emergent marsh and a riverine shrubland, to sample for wetland condition. Both wetlands sampled at the MCWP received ecological condition rankings of a C, indicating moderate deviation from reference condition.

Wetlands at the MCWP are unique in the basin because they were created as part of a restoration effort to provide habitat for migratory waterfowl and shorebirds. Difficulties assessing the ecological condition of created wetlands like these will be further discussed in section 6.2.4.



## 6.2. Upper Basin Vs Floodplain

The northern portion of the study area, referred to herein as the Upper Basin, is located along the western foothills of the Sierra Madre (Figure 14). The landscape consists of upland sagebrush slopes with intermittent drainages that contain wet meadows, willow shrubland wetland, and some beaver-influenced emergent marsh complexes. Cattle grazing is the dominant land use in this area. There is little irrigation and most hydrologic alterations occur in the form of bermed ponds developed in ephemeral drainages to provide water for livestock and wildlife use.

The southern portion of the study area, referred to herein as the Floodplain, lies within the floodplain of the Little Snake River and the lower reaches of Savery Creek. Here the landscape is a mosaic of riparian cottonwood and willow galleries, irrigated pastures and hayfields, and some residential development. Wetlands are associated with the river channel and irrigated areas, and include irrigated wet meadows, riverine oxbows, willow shrublands, and emergent marshes dominated by rushes and cattails.

Land in the Upper Basin is mainly public, whereas the Floodplain is largely privately owned. Approximately 2/3 of the sampled wetlands in the LSRB were located in the Upper Basin (Table 18). More wetlands were sampled in the Upper Basin because we had to sample sites in the order they were selected in our survey design. When a site was rejected from a privately-owned land it was replaced by the next sampleable site on the list. This resulted in more sites being sampled on publicly managed land in the Upper Basin where we did not need permission to sample. It is important to consider the distribution of sampled sites when interpreting our data. The basin-wide results presented in the above sections would look different if more landowners had given permission to same in the Floodplain.

**Table 18.** Number of wetland sites sampled by landowner/manager in the Upper Basin and Floodplain focal areas.

Landowner/manager	Upper Basin	Floodplain
Bureau of Land Management	15	1
State	15	1
Private	16	21
Total	<b>46</b>	<b>23</b>

### 6.2.1 Wetland Types

Wetland subgroups were not equally sampled across focal areas. Over ¾ of the wet meadow and riverine shrubland wetlands sampled were located in the Upper Basin in wetland complexes fed by ephemeral drainages (Table 19). A higher proportion of emergent marsh wetlands were sampled in the Floodplain. Many of these were riverine oxbows located along the Little Snake River or depressional wetlands in old river channels located within irrigated hayfields.



**Table 19.** Number of wetland sites sampled by wetland classification in the Upper Basin and Floodplain focal areas.

Wetland Subgroup	Upper Basin	Floodplain
Emergent Marsh	10	19
Riverine Shrubland	14	4
Wet meadows	22	7
Total	<b>46</b>	<b>23</b>

### 6.2.2 Ecological Condition

The distribution of EIA scores differed between the Upper Basin and the Floodplain focal areas. Specifically, all A-ranked wetlands and most (88%) B-ranked wetlands were located in the Upper Basin. In contrast, 71% of C-ranked wetlands and all D-ranked wetlands (n = 6) were sampled in the Floodplain (Table 20). Ecological condition scores are influenced by alterations to natural hydrology, natural plant communities and/or the impacts from surrounding land use. The below sections help provide context for interpreting EIA scores.

**Table 20.** Number of wetland sites sampled by ecological condition score in the Upper Basin and Floodplain focal areas.

EIA Score	Upper Basin	Floodplain
A	6	-
B	37	5
C	3	12
D	-	6

### 6.2.3 Indicators of Disturbance

Potential indicators of disturbance include natural phenomena or human caused impacts that have the ability to stress a wetland or reduce its ecological condition. By identifying the prevalent disturbances that could impact wetland health in a given area, land managers can address and mitigate impacts from disturbances that are under their control (Lemly and Gilligan 2013).

Indicators of disturbance near or within wetlands sampled in this study reflect differences in land use and potential impacts between the focal areas. For example, we observed many indicators of disturbance associated with agriculture and development in the Floodplain: hay production at 96% of the sampled wetlands, run-off from agricultural practices (observed or inferred) at 87%, irrigation infrastructure at 87%, and residential development at 43% (Table 21). In contrast, wetlands sampled in the Upper Basin had indicators of disturbance from grazing (at 76% of wetlands) and soil degradation by native ungulates and livestock (at 42% of wetlands). Roads

and the presence of non-native species were observed near or within almost all sites sampled in both focal areas.

**Table 21.** Number of wetland sites sampled with disturbance indicators present in the Upper Basin and Floodplain focal areas.

Disturbance Indicator	# of sites		% of Sites	
	Upper Basin	Floodplain	Upper Basin	Floodplain
Agriculture	1	22	2%	96%
Browse	11	4	24%	17%
Residential development	0	10	0%	43%
Filling	1	2	2%	9%
Grazing	34	7	76%	30%
Irrigation infrastructure	11	20	24%	87%
Non-native species	43	23	96%	100%
Agricultural run-off	2	20	4%	87%
Resource extraction	1	2	2%	9%
Roads	39	19	87%	83%
Soil Degradation	19	7	42%	30%
Vegetation conversion	1	2	2%	9%
Water development	13	4	29%	17%

#### 6.2.4 Hydrologic Alteration

Hydrology is the primary driver of the establishment and maintenance of wetlands, affecting the ecological processes that sustain ecosystem function (Maltby and Barker 2009). Therefore, the presence of hydrological alterations (such as irrigation infrastructure) alters the timing and quantity of water available within the basin, directly or indirectly affecting the quantity and type of wetlands present.

The Landscape Hydrology Metric (LHM) is an assessment of alteration to the hydrology of sampled wetlands. LHM uses the degree of human influence on the hydrologic regime to place wetlands into four classes: *historical* wetlands appear to have water sources without human alteration; *created* wetlands, in contrast, appear to depend entirely on water from adjacent irrigated infrastructure ; *supported* wetlands appear to receive water both from natural sources and from irrigation; and *hybrid* wetlands appear to be supported by natural water sources that have been modified to some extent by humans.


Thirty-seven wetlands (80%) in the Upper Basin and only one wetland in the Floodplain were classified as historical based on the LHM analysis (Table 22). Nine wetlands in the Upper Basin

and nine in the Floodplain were identified as hybrid. Seven wetlands in the Floodplain were supported and six wetlands were created. Wetlands supported or created by irrigation infrastructure can be highly vulnerable to changes in water availability (Peck et al. 2004). This is important to consider when prioritizing wetlands for restoration and conservation efforts.

**Table 22.** Number of wetland sites sampled by LHM Classification in the Upper Basin and Floodplain focal areas.

LHM	Upper Basin	Floodplain
Historical	37	1
Hybrid	9	9
Supported	-	7
Created	-	6

Flood irrigation for hay production alters the hydrology of wetlands in the LSRB (WBHCP 2014). Five percent of the entire LSRB (13,374 acre) is mapped as irrigated lands and most of these acres (95%) occur in the Floodplain. Seventy-two percent (5863 acres) of wetland acres in the Floodplain overlap with irrigation (Figure 15). We observed during sampling that more acres are receiving water from irrigation runoff and seepage from irrigation infrastructure than are currently mapped.



**Figure 15.** Spatial distribution of wetland acres, irrigated acres, and wetlands acres overlapping with irrigated acres in the Floodplain focal area.

Highly modified wetlands, like those created or supported by irrigation, or created by restoration efforts such as the MCWP, present a major challenge when assessing wetland condition. The Ecological Integrity Assessment methods used for this survey were designed to evaluate the degree to which wetlands deviate from their natural condition because of hydrologic alterations,

disruptions of native plant communities, and impacts from surrounding land use. EIA heavily weights hydrologic alteration and intensive land use surrounding wetlands negatively in scoring formulas. For this reason, the EIA method fails to recognize the importance of wetlands created or supported by hydrological alterations (irrigation) and restoration efforts. In some instances, it is important to look beyond the low score assigned to wetlands by the EIA method, because they can harbor substantial biodiversity, host multiple species of concern, and can be highly productive systems (Tibbets et al. 2015, Tibbets et al. 2016a, 2016b).

## **7.0 DISCUSSION**

According to the Upper Little Snake River wetland plan (WBHCP 2014), flood irrigation is used on most irrigated lands in the basin. Pressure to change current water management practices has been identified as a moderate threat to wetlands in the LSRB (WBCHP 2014). Water shortages in the Colorado River Basin due to climate alteration and predicted drought (Cook et al. 2004) and increased population (Hansen et al. 2002) are likely to put pressure on Wyoming agricultural producers to alter current irrigation practices and convert to center-pivot irrigation.

Many studies have begun to quantify the importance of irrigation-influenced wetlands for migrating birds and other wildlife (Chester and Robson 2013, Moulton et al. 2013, Patla 2015, Donnelly et al. 2016). In addition to wildlife habitat, there is increased recognition of the ecosystem services provided by irrigation- influenced wetlands (Tanner et al. 2013) for pesticide de-contamination (Tournebize et al. 2013), reduction of nitrogen transport from agricultural catchments, and increases in species richness (Strand and Weisner 2013).

The number and location of wetlands in the LSRB that overlap with irrigation and irrigation infrastructure suggests high vulnerability to wetland loss in the future from conversion to sprinklers or residential development (Copeland et al. 2010, Pocewicz et al. 2014). Approximately 56% of wetland acres in the LSRB overlap with irrigation; conversion to center pivot irrigation could potentially affect an estimated 6,500 acres of wetlands in the basin and the wildlife habitat and ecosystem services they provide. Conservation strategies aimed at protecting lands designated as wetlands may fall short of their intended purpose if water quantity and timing crucial to wetland function are also not retained (Downard and Endter-Wada 2013).

## **8.0 CONCLUSION**

This study provides a broad baseline for beginning to understand the complex relationships between human disturbance, hydrologic alteration, and wetland condition in the LSRB. Additional analyses focusing on the functional and wildlife habitat value of wetlands, as well as the relationship between anthropogenic disturbance and condition are needed. We are just beginning to understand the biodiversity supported by these wetland systems as well as the ecosystem service values that they provide.

In in conjunction with this study, the Nature Conservancy in Wyoming, in collaboration with St. Mary's University of Minnesota, is developing the WyoWet Decision Support Tool. St. Mary's University is updating NWI mapping for 8 USGS Quads along the Little Snake River and the Muddy Creek Wetlands Project which will result in more accurate wetland boundaries. Wetland polygons are being attributed with the Landscape, Landform, Waterbody, Water flow path (LLWW) classification developed by USFWS (Tiner 2003) which can be combined with Cowardin et al. (1979) to estimate functional potential for all wetlands and riparian areas in the LSRB (GeoSpatial Services Saint Mary's University of Minnesota 2018b).

WyoWet allows users to view wetland polygons and interact with associated data that: describes the wetlands biological and hydrologic functional potential; ranks its vulnerability to disturbances; displays hydrologic alterations to the landscape; and displays adjacent landownership/management patterns. WyoWet will give land managers the tools to prioritize restoration, conservation, and protection efforts based on site specific data in the LSRB. The updated mapping and WyoWet tool will be available in 2019.

## LITERATURE CITED

- Adamus, P. 2004. Classification, assessment, monitoring and evaluation of riverine and palustrine wetland ecosystems. *Wetland and Riparian Areas of the Intermountain West: Ecology and Management* 4:277.
- Brinson, M. M. 1993. A Hydrogeomorphic Classification for Wetlands.
- Bureau of Land Management. 2010. Wyoming surface and mineral ownership. Bureau of Land Management.
- Chester, E. T., and B. J. Robson. 2013. Anthropogenic refuges for freshwater biodiversity: their ecological characteristics and management. *Biological Conservation* 166:64–75.
- Comer, P., S. Menard, M. Tuffly, K. Kindscher, R. Rondeau, G. Jones, G. Steinuaer, R. Schneider, and D. Ode. 2003. Upland and Wetland Ecological Systems in Colorado , Wyoming , South Dakota , Nebraska , and Kansas: Report and Map to the National Gap Analysis Program. Page Analysis.
- Cook, E. R., C. A. Woodhouse, C. M. Eakin, D. M. Meko, and D. W. Stahle. 2004. Long-term aridity changes in the western United States. *Science* 306:1015–1018.
- Copeland, H. E., S. A. Tessman, E. H. Girvetz, L. Roberts, C. Enquist, A. Orabona, S. Patla, and J. Kiesecker. 2010a. A geospatial assessment on the distribution, condition, and vulnerability of Wyoming’s wetlands. *Ecological Indicators* 10:869–879.
- Copeland, H., S. A. Tessman, E. H. Girvetz, L. Roberts, C. Enquist, A. Orabona, S. Patla, and J. M. Kiesecker. 2010b. A geospatial assessment on the distribution, condition, and vulnerability of Wyoming’s wetlands. *Ecological Indicators* 10:869–879.
- Costanza, R., R. d’Arge, R. De Groot, S. Farber, M. Grasso, B. Hannon, K. Limburg, S. Naeem, R. V O’neill, and J. Paruelo. 1997. The value of the world’s ecosystem services and natural capital. *nature* 387:253–260.
- Cowardin, L. M., V. Carter, F. C. Golet, and E. T. LaRoe. 1979. Classification of Wetlands and Deepwater Habitats of the United States. *Wildlife Research* 2004:79.
- Culver, D. R., and J. M. Lemly. 2013. Field Guide to Colorado ’ s Wetland Plants: Identification, Ecology, and Conservation. Colorado Natural Heritage Program, Fort Collins, CO.
- Dahl, T. E. 1990. Wetlands losses in the United States 1780’s to 1980’s. *Fish and Wildlife Service* 53:13.
- Deller, A., M. E. Kentula, A. T. Herlihy, A. D. Jacobs, M. E. Kentula, and A. T. Herlihy. 2010. Developing an index of wetland condition from ecological data: An example using HGM functional variables from the Nanticoke watershed, USA. *Ecological Indicators* 10:703–712.
- Donnelly, J. P., D. E. Naugle, C. A. Hagen, and J. D. Maestas. 2016. Public lands and private waters: scarce mesic resources structure land tenure and sage-grouse distributions.

Ecosphere 7:e01208.

- Dorn, R. D. 2001. *Vascular Plants of Wyoming*. 3rd edition. Mountain West Publishing, Cheyenne, WY.
- Downard, R., and J. Endter-Wada. 2013. Keeping wetlands wet in the western United States: Adaptations to drought in agriculture-dominated human-natural systems. *Journal of environmental management* 131:394–406.
- Faber-Langendoen, D., C. Hedge, M. Kost, S. Thomas, L. Smart, R. Smyth, J. Drake, and S. Menard. 2011. Assessment of wetland ecosystem condition across landscape regions: a multi-metric approach. Page Nature Serve. Arlington, VA.
- GeoSpatial Services Saint Mary's University of Minnesota. 2018a. Chain Lakes Wildlife Habitat Management Area NWI data. Wyoming Department of Game and Fish, Cheyenne, Wyoming.
- GeoSpatial Services Saint Mary's University of Minnesota. 2018b. Little Snake River NWI Data. The Wyoming Natural Diversity Database, Laramie, Wyoming.
- Gwin, S. E., M. E. Kentula, and P. W. Shaffer. 1999. Evaluating the effects of wetland regulation through hydrogeomorphic classification and landscape profiles. *Wetlands* 19:477–489.
- Hansen, A., R. Rasker, B. Maxwell, J. Rotella, J. Johnson, A. Parmenter, U. Langner, W. Cohen, R. Lawrence, and M. Kraska. 2002. Ecological Causes and Consequences of Demographic Change in the New West. *BioScience* 52:151–162.
- Johnston, B. C. 2001. Field guide to sedge species of the Rocky Mountain Region: The Genus *Carix* in Colorado, Wyoming, western South Dakota, Western Nebraska, and Western Kansas. U.S. Department of Agriculture, Forest Service, Rocky Mountain Region, Denver, CO.
- Knight, D. H., G. P. Jones, W. A. Reiners, and W. H. Romme. 2014. *Mountains and Plains: The Ecology of Wyoming Landscapes*. 2nd edition.
- Lemly, J., and L. Gilligan. 2012. North Platte River Basin Wetland Profile and Condition Assessment. Colorado Natural Heritage Program, Fort Collins, CO.
- Lemly, J., and L. Gilligan. 2013. Ecological integrity assessment for Colorado wetlands; field manual, version 1.0:1–92.
- Maltby, E., and T. Barker. 2009. *The Wetlands Handbook*, 2 Volume Set. John Wiley & Sons.
- Meade, M. 2010. Public-Private Conservation Efforts in the Little Snake River Basin (WY). Intermountain West Joint Venture.
- Mitsch, W. J., and J. G. Gossilink. 2000. The value of wetlands: Importance of scale and landscape setting. *Ecological Economics* 35:25–33.

- Moulton, C., J. Carlisle, K. Brenner, and R. Cavallaro. 2013. Assessment of Foraging Habitats of Whitefaced Ibis near Two Important Breeding Colonies in Eastern Idaho. Idaho Falls, ID.
- Munsell, A. H. 2013. Munsell Soil Color Charts: With Genuine Munsell Color Chips. Munsell Color.
- Newlon, K., K. M. Ramstead, and H. Jamul. 2013. Southeast Montana wetland assessment: developing and refining Montana's wetland assessment and monitoring strategy. Helena, Montana.
- Olsen, T., and T. Kincaid. 2009. Package 'spsurvey': Spatial Survey Design and Analysis.
- Patla, S. 2015. Upper Green River Wetland Core Complex: Regional Wetland Conservation Plan. Wyoming Wetlands Conservation Strategy, Jackson, WY.
- Peck, D. E., D. M. McLeod, J. P. Hewlett, and J. R. Lovvorn. 2004. Irrigation-dependent wetlands versus instream flow enhancement: economics of water transfers from agriculture to wildlife uses. *Environ Manage* 34:842–855.
- Pocewicz, A., H. E. Copeland, M. B. Grenier, D. A. Keinath, and L. M. Washkoviak. 2014. Vulnerability of Wyoming's. *Terrestrial Wildlife and Habitats*:16.
- R Development Core Team. 2014. R, a Language and Environment for Statistical Computing. R Foundation for Statistical Computing. Vienna, Austria.
- Rocchio, J. 2007. Assessing ecological condition of headwater wetlands in the southern Rocky Mountains using a vegetation Index of biotic integrity:(version 1.0).
- Skinner, Q. D. 2010. A Field Guide to Wyoming Grasses. Education Resources Publishing, Cumming, GA.
- Stevens, D. L., and S. F. Jensen. 2007. Sample design, execution, and analysis for wetland assessment. *Wetlands* 27:515–523.
- Stevens, D. L., A. R. Olsen, D. L. S. T. Jr, and A. R. O. Lsen. 2004. Spatially Balanced Sampling of Natural Resources. *Journal of the American Statistical Association* 99:262–278.
- Stoddard, J. L., A. T. Herlihy, D. V. Peck, R. M. Hughes, T. R. Whittier, and E. Tarquinio. 2008. A process for creating multimetric indices for large-scale aquatic surveys. *Journal of the North American Benthological Society* 27:878–891.
- Stoddard, J. L., D. P. Larsen, C. P. Hawkins, R. K. Johnson, and R. H. Norris. 2006. Setting expectations for the ecological condition of streams: the concept of reference condition. *Ecological Applications* 16:1267–76.
- Strand, J. A., and S. E. B. Weisner. 2013. Effects of wetland construction on nitrogen transport and species richness in the agricultural landscape—Experiences from Sweden. *Ecological Engineering* 56:14–25.



- Swink, F., and G. Wilhelm. 1994. *Plants of the Chicago Region*. 4th Edition. Morton Arboretum, Lisle, IL.
- Swink, F., and G. S. Wilhelm. 1979. *Plants of the Chicago Region*. Revised and expanded edition with keys. The Morton Arboretum, Lisle, IL.
- Tanner, C. C., C. Howard-Williams, M. D. Tomer, and R. Lowrance. 2013. Bringing together science and policy to protect and enhance wetland ecosystem services in agricultural landscapes. *Ecological Engineering* 56:1–4.
- The Wyoming Natural Diversity Database. 2017. The Wyoming Natural Diversity Database Website. <http://www.uwyo.edu/wyndd/>.
- Tibbets, T. M., H. E. Copeland, L. M. Washkoviak, S. Patla, and G. Jones. 2015. *Wetland Profile and Condition Assessment of the Upper Green River Basin, Wyoming*. Lander, Wyoming.
- Tibbets, T. M., L. M. Washkoviak, S. Tessmann, G. Jones, and H. E. Copeland. 2016a. *Wetland Profile and Condition Assessment of the Laramie Plains Wetland Complex , Wyoming*. Lander, Wy.
- Tibbets, T. M., L. M. Washkoviak, S. Tessmann, G. Jones, and H. E. Copeland. 2016b. *Wetland Profile and Condition Assessment of the Goshen Hole Wetland Complex , Wyoming*. Lander, Wy.
- Tiner, R. W. 2003. *Dichotomous Keys and Mapping Codes for Wetland Landscape Position , Landform , Water Flow Path , and Waterbody Type Descriptors:44*.
- Tournebize, J., E. Passeport, C. Chaumont, C. Fesneau, A. Guenne, and B. Vincent. 2013. Pesticide de-contamination of surface waters as a wetland ecosystem service in agricultural landscapes. *Ecological Engineering* 56:51–59.
- U.S. Army Corps of Engineers. 2008. *Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Western Mountains, Valleys, and Cost Regions*.
- U.S. Environmental Protection Agency. 2016. *National Wetland Condition Assessment: 2011 Technical Report*. EPA-843-R-15-006.
- U.S. Fish and Wildlife Service. 1984. *National Wetlands Inventory*. U.S. Department of the Interior, Fish and Wildlife Service, Washington, D.C.
- U.S. Fish and Wildlife Service. 2012. *Partners for Fish and Wildlife Program: Mountain-Prairie Region Strategic Plan, 2012-2016*. Denver, CO:160.
- USDA Farm Service Agency. 2009. *National High Altitude Aerial Photography (NAIP)*. USDA Farm Service Agency.
- Vance, L., K. Newlon, J. Lemly, and G. Jones. 2012. *Assessing the Natural Range of Variability in Minimally Disturbed Wetlands Across the Rocky Mountains: the Rocky Mountain ReMAP*

Project:1–209.

Vasilas, L. M., G. W. Hurt, and C. V Noble. 2010. Field indicators of hydric soils in the United States. United States Department of Agriculture, Natural Resources Conservation Service, in cooperation with the National Technical Committee for Hydric Soils, Washington, DC.

Washkoviak, L. M., B. Heidel, and G. P. Jones. 2017. Floristic Quality Assessment for Wyoming Flora: Developing Coefficients of Conservatism. The Wyoming Natural Diversity Database, Laramie, Wyoming.

Whittier, T. R., S. G. Paulsen, D. P. Larsen, S. A. Peterson, A. T. Herlihy, and P. R. Kaufmann. 2002. Indicators of Ecological Stress and Their Extent in the Population of Northeastern Lakes: A Regional-Scale Assessment. *BioScience* 52:235.

(WBHCP) Wyoming Bird Habitat Conservation Partnership. 2014. Regional Conservation Plan: Upper Little Snake River wetlands Complex.

Wyoming Game and Fish Department. 2010. State Wildlife Action Plan. Cheyenne, WY.

Wyoming Game and Fish Department. 2018. Wyoming Wetlands.  
<https://wgfd.wyo.gov/Habitat/Wyoming-Wetlands>.

Wyoming Joint Ventures Steering Committee. 2010. Wyoming Wetlands Conservation Strategy. Cheyenne, WY.

Wyoming Wildlife Consultants. 2007. Irrigated lands of Wyoming. Wyoming Geographic Information Services (WYGISC), Laramie, Wyoming.

**Appendix A: Field Key to Wetland and Riparian Ecological Systems of Wyoming**  
**Last Updated April 7, 2015**

**1b.** Wetlands and riparian areas of the Western Great Plains. *[If on the edge of the foothills, try both Key A and Key B]* .....  
 ..... **KEY A: WETLANDS AND RIPARIAN AREAS OF THE WESTERN GREAT PLAINS**

**1b.** Wetland and riparian areas west of the Great Plains ..... **2**

**2a.** Wetlands and riparian areas with alkaline or saline soils within the inter-mountains basins of the Rocky Mountains (Upper Green River basin, Wind River basin, etc.) *[If the site does not match any of the descriptions within Key B, try Key C as well. Wetlands and riparian areas of the Rocky Mountains transition into the inter-mountain basins.]*.....  
 ..... **KEY B: WETLANDS AND RIPARIAN AREAS OF THE INTER-MOUNTAIN BASINS**

**2b.** Wetlands and riparian areas of the Rocky Mountains, including the Snowy Mountains, the Wind Rivers, the Absorakas and the Bighorns.. .....  
 ..... **KEY C: WETLANDS AND RIPARIAN AREAS OF THE ROCKY MOUNTAINS**



**Ecological Systems of Wyoming**

- Black Hills
- Inter-mountain Basins
- Rocky Mountains
- Western Great Plains

**KEY A: WETLANDS AND RIPARIAN AREAS OF THE WESTERN GREAT PLAINS**

**1a.** Low stature shrublands dominated by species such as *Sarcobatus vermiculatus*, *Atriplex* spp., *Ericameria nauseosa*, *Artemisia* sp. Vegetation may be sparse and soils may be saline. Sites may be located on the edge alkali depressions, or in flats or washes not typically associated with river and stream floodplains. [These systems were originally described for the Inter-Mountain Basins, but may extend to the plains.] ..... **2**

**1b.** Wetland is not a low stature shrub-dominated saline wash or flat..... **3**

**2a.** Shrublands with sparse (<20%) vegetation cover, located on flats or in temporarily or intermittently flooded drainages, or on the edge of playas and alkali depressions. They are typically dominated by *Sarcobatus vermiculatus* and *Atriplex* spp. with inclusions of *Sporobolus airoides*, *Pascopyrum smithii*, *Distichlis spicata*, *Puccinellia nuttalliana*, and *Eleocharis palustris* herbaceous vegetation ..... **Inter-Mountain Basins Greasewood Flat**

**2b.** Sites with > 20% total vegetation cover and restricted to temporarily or intermittently flooded drainages with a variety of sparse or patchy vegetation including *Sarcobatus vermiculatus*, *Ericameria nauseosa*, *Artemisia* sp., *Grayia spinosa*, *Distichlis spicata*, and *Sporobolus airoides*. ..... **Inter-Mountain Basins Wash**

**3a.** Sites located within the floodplain or immediate riparian zone of a river or stream. Vegetation may be entirely herbaceous or may contain tall stature woody species, such as *Populus* spp. or *Salix* spp. Water levels variable. Woody vegetation that occurs along reservoir edges can also be included here.... **4**

**3b.** Herbaceous wetlands of the Western Great Plains that are isolated or partially isolated from floodplains and riparian zones, often depressional with or without an outlet. .... **8**

**4a.** Herbaceous wetlands within the floodplain with standing water at or above the surface throughout the growing season, except in drought years. Water levels are often high at some point during the growing season, but managed systems may be drawn down at any point depending on water management regimes. Vegetation typically dominated by species of *Typha*, *Scirpus*, *Schoenoplectus*, *Carex*, *Eleocharis*, *Juncus*, and floating genera such as *Potamogeton*, *Sagittaria*, and *Ceratophyllum*. The hydrology may be entirely managed. Water may be brackish or not. Soils are highly variable. This system includes natural warm water sloughs and other natural floodplain marshes as well as a variety of managed wetlands on the floodplain (e.g., recharge ponds, moist soil units, shallow gravel pits, etc.)..... **Western North American Emergent Marsh**

**4b.** Not as above. Wetland and riparian vegetation that typically lacks extensive standing water. Vegetation may be herbaceous or woody. Management regimes variable..... 5

**5a.** Large herbaceous wetlands within the floodplain associated with a high water table that is controlled by artificial overland flow (irrigation). Sites typically lack prolonged standing water. Vegetation is dominated by native or non-native herbaceous species; graminoids have the greatest canopy cover. Species composition may be dominated by non-native hay grasses such as *Poa spp.*, *Alopecurus sp.*, *Phleum pretense*, and *Bromus inermis* spp. *inermis*. There can be patches of emergent marsh vegetation and standing water less than 0.1 ha in size; these are not the predominant vegetation. .... **Irrigated Wet Meadow (not an official Ecological System)**

**5b.** Predominantly natural vegetation (though may be weedy and altered) within the floodplain or immediate riparian zone of a river or stream, dominated by either woody or herbaceous species. Not obviously controlled by irrigation. .... 6

**6a.** Riparian woodlands and shrublands of the Rocky Mountain foothills on the very western margins of the Great Plains. Woodlands are dominated by *Populus* spp. (mainly *Populus angustifolia*.). Common native shrub species include *Salix* spp., *Alnus incana*, *Betula occidentalis*, *Cornus sericea*, and *Crataegus* spp. Sites are most often associated with a stream channel, including ephemeral, intermittent, or perennial streams (Riverine HGM Class). This system can occur on slopes, lakeshores, or around ponds, where the vegetation is associated with groundwater discharge or a subsurface connection to lake or pond water, and may experience overland flow but no channel formation (Slope, Flat, Lacustrine, or Depressional HGM Classes). It is also typically found in backwater channels and other perennially wet but less scoured sites, such as floodplain swales and irrigation ditches. .... **Rocky Mountain Lower Montane-Foothill Riparian Woodland and Shrubland**

**6b.** Riparian woodlands, shrublands and meadows of Wyoming's Western Great Plains. Common native trees are *Populus deltoides*, *Salix amygdaloides*, *Acer negundo*, *Fraxinus pennsylvanica*., and *Ulmus americana*. Common native shrubs include *Salix* spp., *Rosa* spp, and *Symphoricarpos* spp. Common non-native trees and shrubs are *Tamarix* spp. and *Elaeagnus angustifolia*. .... 7

**7a.** Riparian woodlands, shrublands, and meadows along medium and small rivers and streams. Sites have less floodplain development and flashier hydrology than the next, and all streamflow may drawdown completely for some portion of the year. Water sources include snowmelt runoff (more common in Wyoming), groundwater (prairie streams), and summer rainfall. Dominant species include *Populus deltoides*, *Salix* spp., *Fraxinus pennsylvanica*, *Pascopyrum smithii*, *Panicum* sp., *Carex* spp., *Tamarix* spp., *Elaeagnus angustifolia*, and other non-native grasses and forbs.....  
 .....**Western Great Plains Riparian**

**7b.** Woodlands, shrublands, and meadows along large rivers (the North Platte and its larger tributaries) with extensive floodplain development and periodic flooding that is more associated with snowmelt and seasonal dynamics in the mountains than with local precipitation events. Hydroperiod alterations from major dams and reservoirs alter historic flooding patterns. Dominant communities within this system range from floodplain forests to wet meadow patches, to gravel/sand flats dominated by early successional herbs and annuals; however, they are linked by underlying soils and the flooding regime. Dominant species include *Populus deltoides* and *Salix* spp., *Panicum* sp. and *Carex* spp. *Tamarix* spp., *Elaeagnus angustifolia*, and non-native grasses.....**Western Great Plains Floodplain**

**8a.** Natural shallow depressional wetlands in the Western Great Plains with an impermeable soil layer, such as dense hardpan clay that causes periodic ponding after heavy rains. Sites generally have closed contour topography and are surrounded by upland vegetation. Hydrology is typically tied to precipitation and runoff but lacks a groundwater connection; however some of these sites are receiving increased water from irrigation seepage. Ponding is often ephemeral and sites may be dry throughout the entire growing season during dry years. Species composition depends on soil salinity, may fluctuate depending on seasonal moisture availability, and many persistent species may be upland species. *[The wetlands within this group are collectively referred to **playas or playa lakes**. Ecological systems listed below separate playas based on the level of salinity and total cover of vegetation.]*..... **9**

**8b.** Herbaceous wetlands in the Western Great Plains not associated with hardpan clay soils. Sites may or may not be depressional and may or may not be natural. .... **10**

**9a.** Shallow depressional wetlands with less saline soils than the next. Dominant species are typically not salt-tolerant. Sites may have obvious vegetation zonation of tied to water levels, with the most hydrophytic species occurring in the wetland center where ponding lasts the longest. Common native species include *Pascopyrum smithii*, *Iva axillaris*, , *Eleocharis* spp., *Oenothera canescens*, *Plantago* spp., *Polygonum* spp., *Conyza canadensis* ,and *Phyla cuneifolia*. Non-native species are very common in these sites, including *Salsola australis*, *Bassia sieversiana*, *Verbena bracteata*, and *Polygonum aviculare*. Sites have often been affected by agriculture and heavy grazing. Many have been dug out or “pitted” to increase water retention and to tap shallow groundwater.....  
 .....**Western Great Plains Closed Depression Wetland**

**9b.** Shallow depressional herbaceous wetlands with saline soils. Salt encrustations can occur on the surface. Species are typically salt-tolerant, including *Distichlis spicata*, *Puccinellia nuttalliana*, *Salicornia rubra*, *Schoenoplectus maritimus*, *Schoenoplectus americanus*, *Suaeda calceoliformis*, *Spartina* spp., *Triglochin maritima*, and occasional shrubs such as *Sarcobatus vermiculatus*. [This system resembles the Inter-Mountain Basins Alkaline Closed Depression but occur in the Great Plains ecoregion. Note: Low stature shrub-dominant wetlands key in the flats and wash systems above.].....  
 ..... **Western Great Plains Saline Depression Wetland**

**10a.** Herbaceous wetlands with standing water at or above the surface throughout the growing season, except in drought years. Water levels are often high at some point during the growing season, but managed systems may be drawn down at any point depending on water management regimes. Vegetation typically dominated by species of *Typha*, *Scirpus*, *Schoenoplectus*, *Carex*, *Eleocharis*, *Juncus*, and floating genera such as *Potamogeton*, *Sagittaria*, and *Ceratophyllum*. The isolated expression of this system can occur around ponds, as fringes around lakes, and at any impoundment of water, including irrigation run-off. The hydrology may be entirely managed or artificial. Water may be brackish or not. Soils are highly variable..... **Western North American Emergent Marsh**

**10b.** Herbaceous wetlands associated with a high water table that is controlled by artificial overland flow (irrigation) or artificial groundwater seepage (including from leaky irrigation ditches). Sites typically lack prolonged standing water. Vegetation is dominated by native or non-native herbaceous species; graminoids have the greatest canopy cover. Patches of emergent marsh vegetation and standing water are less than 0.1 ha in size and not the predominant vegetation.....  
 ..... **Irrigated Wet Meadow (not an official Ecological System)**

**KEY B: WETLANDS AND RIPARIAN AREAS OF THE INTER-MOUNTAIN BASINS**

**1a.** Depressional, herbaceous wetlands occurring within dune fields of the inter-mountain basins (e.g. Great Divide basin)..... **Inter-Mountain Basins Interdunal Swale Wetland**

**1b.** Wetlands not associated with dune fields ..... **2**

**2a.** Depressional wetlands. Soils are typically alkaline to saline clay with hardpans. Salt encrustation typically visible on the soil surface or along the water edge. Water levels various. Cover of vegetation variable, can be extremely sparse (<10% cover) or moderate to high (30–60% cover). Typically herbaceous dominated, but may contain salt-tolerant shrubs on the margins..... **3**

**2b.** Non-depressional wetlands on flats or in washes, with alkaline to saline soils. Cover of vegetation variable, can be extremely sparse (<10% cover) or moderate to high (30–60% cover). Typically shrub dominated. Most common species are *Sarcobatus vermiculatus* and *Atriplex* spp..... **4**



**3a.** Depressional, alkaline wetlands that are seasonally to semi-permanently flooded, usually retaining water into the growing season and drying completely only in drought years. Many are associated with irrigation seepage, springs, or located in large basins with internal drainage. Seasonal drying exposes mudflats colonized by annual wetland vegetation. This system can occur in alkaline basins and swales and along the drawdown zones of lakes and ponds. They generally have thick unvegetated salt crusts over clay soils surrounded by zones of vegetation transitioning to the uplands. In these zones vegetation cover is generally >10% and species are typically salt-tolerant such as *Distichlis spicata*, *Puccinellia* spp., *Leymus* sp., *Schoenoplectus maritimus*, *Schoenoplectus americanus*, *Triglochin maritima*, and *Salicornia* spp.....**Inter-Mountain Basins Alkaline Closed Depression**

**3b.** Barren and sparsely vegetated playas (generally <10% plant cover. Could be more if annuals or upland vegetation are encroaching). Salt crusts are common throughout, with small saltgrass beds in depressions and sparse shrubs around the margins. These systems are intermittently flooded. The water generally comes from precipitation and is prevented from percolating through the soil by an impermeable soil sub horizon and is left to evaporate. Soil salinity varies with soil moisture and greatly affects species composition. Characteristic species may include *Sarcobatus vermiculatus*, *Distichlis spicata*, and/or *Atriplex* spp..... **Inter-Mountain Basins Playa**

**4a.** Shrublands with >10% total vegetation cover, located on flats. Vegetation dominated by *Sarcobatus vermiculatus* and *Atriplex* spp. with inclusions of *Artemisia tridentata* ssp. *Tridentata*, *Sporobolus airoides*, *Pascopyrum smithii*, *Distichlis spicata*, *Puccinellia nuttalliana*, and herbaceous vegetation.....  
..... **Inter-Mountain Basins Greasewood Flat**

**4b.** Sites with < 10% total vegetation cover and restricted to temporarily or intermittently flooded drainages with a variety of sparse or patchy vegetation including *Sarcobatus vermiculatus*, *Ericameria nauseosa*, *Artemisia cana*, *Artemisia tridentata*, *Distichlis spicata*, and *Sporobolus airoides*. .....  
..... **Inter-Mountain Basins Wash**

**KEY C: WETLANDS AND RIPARIAN AREAS OF THE ROCKY MOUNTAINS**

**1a.** Wetland defined by groundwater inflows and organic soil (peat) accumulation of at least 40 cm in the upper 80 cm. Vegetation can be woody or herbaceous. If the wetland occurs within a mosaic of non-peat forming wetland or riparian systems, then the patch must be at least 0.1 hectare (0.25 acre). If the wetland occurs as an isolated patch surrounded by upland, then there is no minimum size criterion. ....  
..... **Rocky Mountain Subalpine-Montane Fen**

**1b.** Wetland does not have at least 40 cm of organic soil (peat) accumulation or occupies an area less than 0.1 hectares (0.25 acres) within a mosaic of other non-peat forming wetland or riparian systems... 2

**2a.** Total woody canopy cover generally 25% or more within the overall wetland/riparian area. Any purely herbaceous patches are less than 0.5 hectare and occur within a matrix of woody vegetation. [Note: Relictual woody vegetation such as standing dead trees and shrubs are included here.] ..... 3

**2b.** Total woody canopy cover generally less than 25% within the overall wetland/riparian area. Any woody vegetation patches are less than 0.5 hectare and occur within a matrix of herbaceous wetland vegetation ..... 5

**3a.** Riparian woodlands and shrublands of the foothill and lower montane zones on the Rocky Mountains. Woodlands are dominated by *Populus* spp. (*Populus angustifolia*, or the hybrid *P. acuminata*). At higher elevations *Picea engelmannii*, *Abies lasiocarpa*, *Pseudotsuga menziesii*, and *Pinus ponderosa* can be found. Common native shrub species include *Salix* spp., *Alnus incana*, *Betula occidentalis*, *Cornus sericea*, and *Crataegus* spp. Sites are most often associated with a stream channel, including ephemeral, intermittent, or perennial streams (Riverine HGM Class). This system can occur on slopes, lakeshores, or around ponds, where the vegetation is associated with groundwater discharge or a subsurface connection to lake or pond water, and may experience overland flow but no channel formation (Slope, Flat, Lacustrine, or Depressional HGM Classes). It is also typically found in backwater channels and other perennially wet but less scoured sites, such as floodplain swales and irrigation ditches. (this system is also found in the inter-mountain basin ecoregion).. .....  
.....**Rocky Mountain Lower Montane-Foothill Riparian Woodland and Shrubland**

**3b.** Riparian woodlands and shrublands of the montane or subalpine zone ..... 4

**4a.** Montane or subalpine riparian woodlands (canopy dominated by trees). This system occurs as a narrow streamside forest lining small, confined low- to mid-order streams. Common tree species include *Abies lasiocarpa*, *Picea engelmannii*, and *Populus tremuloides* (The overstory consists of *Picea engelmannii*, often with some *Abies lasiocarpa* and *Populus tremuloides*. These riparian areas generally occur at elevations where the uplands support upper montane and subalpine forests -- *Pinus contorta*, *Picea engelmannii*, *Abies lasiocarpa*. The common riparian trees in this type -- *Picea engelmannii*, *Abies lasiocarpa*, *Populus tremuloides* -- also grow in riparian zones in the lower montane, but there they are joined by *Populus angustifolia*, sometimes *Populus acuminata*, *Populus balsamifera* (mostly in NW Wyoming), *Picea pungens* (NW Wyoming : Snake River drainage, and the Wind River around Dubois), *Pseudotsuga menziesii*, *Pinus ponderosa* (eastern half of WY). Then, with decreasing elevation, the conifer drop out, *Populus acuminata* increases, and *Populus deltoides* becomes a major species.) .....  
..... **Rocky Mountain Subalpine-Montane Riparian Woodland**

**4b.** Montane or subalpine shrub wetlands (canopy dominated by shrubs with sparse or no tree cover). This system is most often associated with streams (Riverine HGM Class), occurring as either a narrow band of shrubs lining streambanks of steep V-shaped canyons (straight, with boulder and cobble substrate) or as a wide, extensive shrub stand on alluvial terraces in low-gradient valley bottoms (more sinuous, with finer-textured substrates. Sometimes referred to as a *shrub carr*). Beaver activity is common within the wider occurrences. In addition, this system can occur around the edges of fens, lakes, seeps, and springs on slopes away from valley bottoms. This system can also occur within a mosaic of multiple shrub- and herb-dominated communities within snowmelt-fed basins. In all cases, vegetation is dominated by species of *Salix*, *Alnus*, or *Betula* but their composition varies depending on stream gradient. *Alnus incana* is a dominant or co-dominant along high-gradient streams; *Betula occidentalis* often co-dominates. Willows are present, as is *Cornus sericea*, but rarely dominate. In contrast, along the lower-gradient streams in wide valleys, the willows dominate; *Betula* and *Cornus* often are present but secondary to the willows; *Alnus* usually is a minor component. . . . .

..... **Rocky Mountain Subalpine-Montane Riparian Shrubland**

**5a.** Herbaceous wetlands with water present throughout all or most of the year. Water is at or above the surface throughout the growing season, except in drought years. This system can occur around ponds, as fringes around lakes, and along slow-moving streams and rivers. The vegetation is dominated by common emergent and floating leaved plants, including species of *Scirpus*, *Schoenoplectus*, *Typha*, *Juncus*, *Carex*, *Potamogeton*, *Polygonum*, and *Nuphar*.....

..... **Western North American Emergent Marsh**

**5b.** Herbaceous wetlands that typically lack extensive standing water. Patches of emergent marsh vegetation and standing water are less than 0.1 ha in size and not the predominant vegetation. .... **7**

**6a.** Herbaceous wetlands associated with a high water table or overland flow, but typically lack standing water. Sites with *no channel formation* are typically associated with snowmelt or groundwater and not subjected to high disturbance events such as flooding (Slope HGM Class). Sites *associated with a stream channel* are more tightly connected to overbank flooding from the stream channel than with snowmelt and groundwater discharge. Vegetation is dominated by herbaceous species; typically graminoids have the highest canopy cover including *Carex* spp., *Calamagrostis* spp., and *Deschampsia caespitosa*.....

..... **Rocky Mountain Alpine-Montane Wet Meadow**

**6b.** Large herbaceous wetlands associated with a high water table that is controlled by artificial overland flow (irrigation). Sites typically lack prolonged standing water, but may have standing water early in the season if water levels are very high. Vegetation is dominated by native or non-native herbaceous species; graminoids have the highest canopy cover .....

..... **Irrigated Wet Meadow (not an official Ecological System)**

## **Appendix B. Operational Definition of a Wetland for Condition Assessment of the Little Snake River Basin, Wyoming**

The operational definition of wetlands used in this project is based on the definition adopted by the U.S. Fish and Wildlife Service (USFWS) and used in the National Wetland Inventory (Cowardin et al. 1979):

*“Wetlands are lands transitional between terrestrial and aquatic systems where the water table is usually at or near the surface or the land is covered by shallow water. For purposes of this classification wetlands must have one or more of the following attributes: (1) at least periodically, the land supports predominantly hydrophytes; (2) the substrate is predominantly undrained hydric soil; and (3) the substrate is nonsoil and is saturated with water or covered by shallow water at some time during the growing season of each year.”*

However, it is important to note that standard wetland delineation techniques are based on a different definition used by the U.S. Army Corps of Engineers (ACOE) and the Environmental Protection Agency (EPA) for regulatory purposes under Section 404 of the Federal Clean Water Act:

*“[Wetlands are] those areas that are inundated or saturated by surface or ground water at a frequency and duration sufficient to support, and under normal circumstances do support, a prevalence of vegetation typically adapted for life in saturated soil conditions.”*

The primary difference between the two definitions is the ACOE/EPA definition requires positive identification of all three wetland parameters (hydrology, vegetation, and soils), whereas the USFWS definition requires only one characteristic must be present. We required two wetland parameters to be present to qualify for assessment sampling. Deep water habitats that would be considered wetlands under the USFWS definition were excluded from this study.

## Appendix C. Little Snake Basin Wetland Assessment Field Form

LOCATION AND GENERAL INFORMATION	
Point Code _____ Date: _____ Surveyors: _____	
Access Comments (note permit requirement or difficulties accessing the site):  	
GPS COORDINATES OF TARGET POINT AND ASSESSMENT AREA	
<u>Dimensions of AA:</u> ___ 40 m radius circle ___ Rectangle ___ Freeform, describe and take a GPS Track	Elevation (m): _____ Target Wetland: ___ Yes ___ No / Type: if no what is new target type: _____ Relation to AA: ___ Centered ___ Included ___ Outside
AA-Center WP #: _____ LAT: _____ LONG: _____ Error (+/-): _____ (Circle AAs Only)	
AA-Track Track Name: _____ Area: _____	
PHOTOS OF ASSESSMENT AREA (Taken at the center point looking out in the cardinal directions for standard 40 m radius circle AA's or from four points on edge of AA looking in for freeform AA's)	
AA-1	WP/Photo #: _____ Aspect: _____ LAT: _____ LONG: _____
AA-2	WP/Photo #: _____ Aspect: _____ LAT: _____ LONG: _____
AA-3	WP/Photo #: _____ Aspect: _____ LAT: _____ LONG: _____
AA-4	WP/Photo #: _____ Aspect: _____ LAT: _____ LONG: _____
Additional AA Photos and Comments:  (Note range of photo numbers and explain particular photos of interest)	
ENVIRONMENTAL DESCRIPTION AND CLASSIFICATION OF ASSESSMENT AREA	
<u>Non-target Inclusions:</u> % AA with > 1m standing water: _____ % AA with Non-target inclusions: _____ Non-target description: _____	<u>Wetland origin (if known):</u> ___ Natural feature with minimal alteration ___ Natural feature, but altered or augmented by modification ___ Non-natural feature created by passive or active management ___ Unknown
<u>Ecological System:</u> (see manual for key and rules on inclusions and pick the <i>best match</i> ) Fidelity: High Med Low	

**ENVIRONMENTAL DESCRIPTION AND CLASSIFICATION OF ASSESSMENT AREA (continued)**

Cowardin Classification (pick one each that best represents AA)

Fidelity: High Med Low

(see manual and pick one each of System, Class, Water Regime, and optional Modifier for dominant type)

HGM Class (pick only one that best represents AA)

Fidelity: High Med Low

\_\_\_ Riverine\*                      \_\_\_ Lacustrine Fringe

\_\_\_ Depressional                \_\_\_ Slope

\_\_\_ Flats                          \_\_\_ Irrigated (choose additional class)

*\*Specific classification and metrics apply to the Riverine HGM Class*

**AA REPRESENTATIVENESS**

Is AA the entire wetland? \_\_\_ Yes \_\_\_ No      If no, is AA representative of larger wetland? \_\_\_ Yes \_\_\_ No

Provide comments:

**Wildlife Observation – record any wildlife observations from site. List species of and type of observation**

Species:	# individuals	Nests	Visible	Vocal	Tracks	Scat

Wildlife Comments:

**ASSESSMENT AREA DRAWING AND COMMENTS**

Add north arrow and approx. scale bar. Document **Community types and abiotic zones** (particularly open water), inflows and outflows, and indicate direction of drainage. Include sketch of soil pit placement. If appropriate, add a **cross-sectional diagram** and indicate slope of side.

**ASSESSMENT AREA SETTING AND SURROUNDING LANDSCAPE DESCRIPTION**

Overall site description and details on site hydrology, soil, and vegetation. Include general landscape setting, dominant plants in buffer, and information on any target wetland types occurring with AA.



AA GROUND COVER AND VERTICAL STRATA					
<b>Cover Classes 1: trace 2: &lt;1% 3: 1–&lt;2% 4: 2–&lt;5% 5: 5–&lt;10% 6: 10–&lt;25% 7: 25–&lt;50% 8: 50–&lt;75% 9: 75–&lt;95% 10: &gt;95% (Unless otherwise noted)</b>					
Cover of standing water of any depth, vegetated or not:					
Cover of running water of any depth, vegetated or not:					
Cover of open water (plant canopy cover < 10%)					
Cover of water with emergent vegetation:					
Cover of water with floating or submerged vegetation:					
Cover of exposed bare ground* – soil / sand / sediment					
Cover of exposed bare ground* – gravel / cobble (~2–250 mm)					
Cover of exposed bare ground* – bedrock / rock / boulder (>250 mm)					
Cover salt crust (all cover, <u>including over vegetation or litter cover</u> )					
Cover of litter (all cover, <u>including under water or vegetation</u> )					
Depth of litter (cm) – average of four non-trampled locations where litter occurs Depth 1 _____ cm    Depth 2 _____ cm    Depth 3 _____ cm    Depth 4 _____ cm    Ave depth:					
Predominant litter type (C = coniferous, E = broadleaf evergreen, D = deciduous, S = sod/thatch, F = forb)					
Cover of standing dead trees (>5 cm diameter at breast height)					
Cover of standing dead shrubs or small trees (<5 cm diameter at breast height)					
Cover of downed coarse woody debris (fallen trees, rotting logs, >5 cm diameter)					
Cover of downed fine woody debris (<5 cm diameter)					
Cover bryophytes (all cover, <u>including under water, vegetation or litter cover</u> )					
Cover lichens (all cover, <u>including under water, vegetation or litter cover</u> )					
Cover algae (all cover, <u>including under water, vegetation or litter cover</u> )					
<b>Height Classes 1:&lt;0.5 m 2: 0.5–1m 3: 1–2 m 4: 2–5 m 5: 5–10 m 6: 10–15 m 7: 15–20 m 8: 20–35 m 9: 35–50 m 10:&gt;50 m</b>					
Vertical Vegetation Strata(live or very recently dead)			Cover / Height →	C	H
(T1)	Dominant canopy trees (>5 m and > 30% cover)				
(T2)	Sub-canopy trees (> 5m but < dominant canopy height) or trees with sparse cover				
(S1)	Canopy layer 2–5 m includes both Tall shrubs or older tree saplings				
	Older tree saplings 2–5 m				
	Tall shrubs 2–5 m				
(S2)	Canopy layer 0.5 – 2 includes both Short shrubs or young tree saplings (0.5–2 m)				
	Young tree saplings 0.5–2 m				
	Short shrubs 0.5–2 m				
(S3)	Dwarf shrubs or tree seedlings (<0.5 m; included short <i>Vaccinium</i> spp., etc.)				
	Tree seedlings <0.5 m				
	Dwarf shrubs <0.5 m (included short <i>Vaccinium</i> spp., etc.)				
(HT)	Herbaceous total				
(H1)	Graminoids (grass and grass-like plants)				
(H2)	Forbs (all non-graminoids)				
(H3)	Ferns and fern allies				
(AQ)	Submergent or floating aquatics				

## Vegetation Species List

Walk through the AA and identify as many plant species as possible beginning with the most dominant species first. Spend *no more* than 1 hour compiling the species list. Once the species list is compiled, estimate absolute cover for each species.

Scientific Name or Pseudonym	% Cover	Coll #	Photos



**SOIL PROFILE DESCRIPTION – SOIL PIT**     **Representative Pit?**

**GPS Waypoint** \_\_\_\_\_    Lat: \_\_\_\_\_    Long: \_\_\_\_\_    Temp \_\_\_\_\_    pH \_\_\_\_\_    EC \_\_\_\_\_    TDS \_\_\_\_\_    Salinity \_\_\_\_\_

Settling Time: \_\_\_\_\_    Depth to saturated soil (cm): \_\_\_\_\_    Depth to free water (cm): \_\_\_\_\_     Not observed, if so:     Pit is filling slowly    OR     Pit appears dry

Depth (cm)	<u>Matrix</u>	<u>Dominant Redox Features</u>		<u>Secondary Redox Features</u>		Texture	% Roots	% Gravel	Remarks
	Color (moist)	Color (moist)	%	Color (moist)	%				
_____	_____	_____	_____	_____	_____	_____	_____	_____	_____
_____	_____	_____	_____	_____	_____	_____	_____	_____	_____
_____	_____	_____	_____	_____	_____	_____	_____	_____	_____
_____	_____	_____	_____	_____	_____	_____	_____	_____	_____
_____	_____	_____	_____	_____	_____	_____	_____	_____	_____

**Hydric Soil Indicators:** See field manual for descriptions and check all that apply to pit.

Histosol (A1)                       Gleyed Matrix (S4/F2)                       Surface Salt Crusts  
 Histic Epipedon (A2/A3)                       Depleted Matrix (A11/A12/F3)                       Translocated Salts  
 Mucky Mineral (S1/F1)                       Redox Concentrations (S5/F6/F8)  
 Hydrogen Sulfide Odor (A4)                       Redox Depletions (S6/F7)

Comments: \_\_\_\_\_

**SOIL PROFILE DESCRIPTION – SOIL PIT 2**     **Representative Pit?**

**GPS Waypoint** \_\_\_\_\_    Lat: \_\_\_\_\_    Long: \_\_\_\_\_    Temp \_\_\_\_\_    pH \_\_\_\_\_    EC \_\_\_\_\_    TDS \_\_\_\_\_    Salinity \_\_\_\_\_

Settling Time: \_\_\_\_\_    Depth to saturated soil (cm): \_\_\_\_\_    Depth to free water (cm): \_\_\_\_\_     Not observed, if so:     Pit is filling slowly    OR     Pit appears dry

Depth (cm)	<u>Matrix</u>	<u>Dominant Redox Features</u>		<u>Secondary Redox Features</u>		Texture	% Roots	% Gravel	Remarks
	Color (moist)	Color (moist)	%	Color (moist)	%				
_____	_____	_____	_____	_____	_____	_____	_____	_____	_____
_____	_____	_____	_____	_____	_____	_____	_____	_____	_____
_____	_____	_____	_____	_____	_____	_____	_____	_____	_____
_____	_____	_____	_____	_____	_____	_____	_____	_____	_____
_____	_____	_____	_____	_____	_____	_____	_____	_____	_____

**Hydric Soil Indicators:** See field manual for descriptions and check all that apply to pit.

Histosol (A1)                       Gleyed Matrix (S4/F2)                       Surface Salt Crusts  
 Histic Epipedon (A2/A3)                       Depleted Matrix (A11/A12/F3)                       Translocated Salts  
 Mucky Mineral (S1/F1)                       Redox Concentrations (S5/F6/F8)  
 Hydrogen Sulfide Odor (A4)                       Redox Depletions (S6/F7)

Comments: \_\_\_\_\_

**SOIL PROFILE DESCRIPTION – SOIL PIT 3**     Representative Pit

GPS Waypoint \_\_\_\_\_ Lat: \_\_\_\_\_ Long: \_\_\_\_\_ Temp \_\_\_\_\_ pH \_\_\_\_\_ EC \_\_\_\_\_ TDS \_\_\_\_\_ Salinity \_\_\_\_\_

Settling Time: \_\_\_\_\_ Depth to saturated soil (cm): \_\_\_\_\_ Depth to free water (cm): \_\_\_\_\_  Not observed, if so:  Pit is filling slowly OR  Pit appears dry

Depth (cm)	Matrix	Dominant Redox Features		Secondary Redox Features		Texture	% Roots	% Gravel	Remarks
	Color (moist)	Color (moist)	%	Color (moist)	%				
_____	_____	_____	_____	_____	_____	_____	_____	_____	_____
_____	_____	_____	_____	_____	_____	_____	_____	_____	_____
_____	_____	_____	_____	_____	_____	_____	_____	_____	_____
_____	_____	_____	_____	_____	_____	_____	_____	_____	_____
_____	_____	_____	_____	_____	_____	_____	_____	_____	_____

**Hydric Soil Indicators:** See field manual for descriptions and check all that apply to pit.

<input type="checkbox"/> Histosol (A1)	<input type="checkbox"/> Gleyed Matrix (S4/F2)	<input type="checkbox"/> Surface Salt Crusts
<input type="checkbox"/> HisticEpipedon (A2/A3)	<input type="checkbox"/> Depleted Matrix (A11/A12/F3)	<input type="checkbox"/> Translocated Salts
<input type="checkbox"/> Mucky Mineral (S1/F1)	<input type="checkbox"/> Redox Concentrations (S5/F6/F8)	
<input type="checkbox"/> Hydrogen Sulfide Odor (A4)	<input type="checkbox"/> Redox Depletions (S6/F7)	

Comments: \_\_\_\_\_

**WATER QUALITY**

Site 1: GPS Waypoint \_\_\_\_\_ Lat: \_\_\_\_\_ Long: \_\_\_\_\_ **Standing OR Flowing**

Temp \_\_\_\_\_ pH \_\_\_\_\_ EC \_\_\_\_\_ TDS \_\_\_\_\_ Salinity \_\_\_\_\_

Site 2: GPS Waypoint \_\_\_\_\_ Lat: \_\_\_\_\_ Long: \_\_\_\_\_ **Standing OR Flowing**

Temp \_\_\_\_\_ pH \_\_\_\_\_ EC \_\_\_\_\_ TDS \_\_\_\_\_ Salinity \_\_\_\_\_

Water quality measurement comments: \_\_\_\_\_

\*Be sure to mark down any soils and water chemistry units

Macro Invertebrate sample taken: _____	Macro invertebrate comments: _____
--	------------------------------------

## LEVEL 2 ECOLOGICAL INTEGRITY ASSESSMENT

### 1. LANDSCAPE CONTEXT METRICS – Check the applicable box.

1a. PERCENT NATURAL LAND COVER				
Select the statement that best describes the <b>percent of natural land cover</b> within 100, 200 and 500 m envelopes surrounding the AA. To determine, identify any patches of natural land cover within the 100, 200 and 500 m envelopes and estimate their total percent of the envelopes. See definitions in the field manual of natural land cover types. Natural land cover patches <i>do not</i> need to be contiguous with the AA.	Distance from AA:	<b>100m</b>	<b>200 m</b>	<b>500 m</b>
	Intact: Landscape contains 90–100% natural land cover.			
	Variegated: Landscape contains 60–90% natural land cover.			
	Fragmented: Landscape contains 20–60% natural land cover.			
	Relictual: Landscape contains <20% natural land cover.			
1b. LANDSCAPE FRAGMENTATION				
Select the statement that best describes the <b>landscape fragmentation</b> with in a 500 m envelope surrounding the AA. To determine, identify the largest unfragmented block <i>that includes the AA</i> within the 500 m envelope and estimate its percent of the total envelope. Well-traveled dirt roads and major canals count as fragmentation, but hiking trails, hayfields, fences and small ditches can be included in unfragmented blocks (see definitions).	Intact: AA embedded in >90–100% unfragmented, natural landscape.			
	Variegated: AA embedded in >60–90% unfragmented, natural landscape.			
	Fragmented: AA embedded in >20–60% unfragmented, natural landscape.			
	Relictual: AA embedded in ≤20% unfragmented, natural landscape.			
1b. RIPARIAN CORRIDOR CONTINUITY (RIVERINE WETLANDS ONLY)				
<i>For riverine wetlands</i> , select the statement that best describes the <b>riparian corridor continuity</b> within 500 m upstream and downstream of the AA. To determine, identify any non-buffer patches (see definitions) within the potential riparian corridor (natural geomorphic floodplain) both upstream and downstream of the AA. Estimate the percentage of the riparian corridor they occupy. <i>For AAs on one side of a very large river channel (~20 m width)</i> , only consider the riparian corridor on that side of the channel.	Intact: >95–100% natural habitat within the riparian corridor both upstream and downstream.			
	Variegated: >80–95% natural within the riparian corridor both upstream and downstream.			
	Fragmented: >50–80% natural habitat within the riparian corridor both upstream and downstream.			
	Relictual: ≤50% natural habitat within the riparian corridor both upstream and downstream.			
Landscape fragmentation and riparian corridor continuity comments:				

1c. BUFFER EXTENT		
Select the statement that best describes the <b>extent of buffer land cover</b> surrounding the AA. To determine, estimate the percent of the AA surrounded by buffer land covers (see definitions). Each segment must be $\geq 5$ m wide and extend along $\geq 10$ m of the AA perimeter.	Buffer land covers surround $>100\%$ of the AA.	
	Buffer land covers surround $>75\text{--}100\%$ of the AA.	
	Buffer land covers surround $>50\text{--}75\%$ of the AA.	
	Buffer land covers surround $>25\text{--}50\%$ of the AA.	
	Buffer land covers surround $\leq 25\%$ of the AA.	
1d. BUFFER WIDTH		
Select the statement that best describes the <b>buffer width</b> . To determine, estimate width (up to 200 m from AA) along eight lines radiating out from the AA at the cardinal and ordinal directions (N, NE, E, SE, S, SW, W, NW).		
1: _____	5: _____	Average buffer width is $>200$ m
2: _____	6: _____	Average buffer width is $>100\text{--}200$ m
3: _____	7: _____	Average buffer width is $>50\text{--}100$ m
4: _____	8: _____	Average buffer width is $>25\text{--}50$ m
Average width: _____		Average buffer width is $\leq 25$ m OR no buffer exists
1e. BUFFER CONDITION		
Select the statement that best describes the <b>buffer condition</b> . Select one statement per column. Only consider <u>the actual buffer</u> measured in metrics 1c and 1d. Use the Landscape Stressor list below to help inform your buffer condition decision		
Abundant ( $\geq 95\%$ ) relative cover native vegetation and little or no ( $<5\%$ ) cover of non-native plants.		Intact soils, little or no trash or refuse, and no evidence of human visitation. Light grazing can be present.
Substantial ( $\geq 75\text{--}95\%$ ) relative cover of native vegetation and low ( $5\text{--}25\%$ ) cover of non-native plants.		Intact or moderately disrupted soils, moderate or lesser amounts of trash, light grazing to moderate grazing OR minor intensity of human visitation or recreation
Moderate ( $\geq 50\text{--}75\%$ ) relative cover of native vegetation.		Moderate or extensive soil disruption, moderate or greater amounts of trash, moderate to heavy grazing OR moderate intensity of human use.
Low ( $<50\%$ ) relative cover of native vegetation OR no buffer exists.		Barren ground and highly compacted or otherwise disrupted soils, moderate or greater amounts of trash, moderate or greater intensity of human use, very heavy grazing OR no buffer exists.
Buffer comments:		



**LANDSCAPE STRESSORS**

Using the table below, identify all **landscape stressor / land uses within a 200 and 500 m envelope** of the AA. Stressors can overlap (e.g., Grazing and moderate recreation can both be counted in the same portion of the envelope). Rank the top 3 stressors effecting the wetland within the 200m and 500m buffers.

<i>Landscape stressor/ land use categories</i>	200m		500m	
	<i>Present</i>	<i>Rank</i>	<i>Present</i>	<i>Rank</i>
Paved roads, parking lots, railroad tracks				
Unpaved roads (e.g., driveway, tractor trail, 4-wheel drive roads)				
Domestic or commercially developed buildings				
Trash or refuse				
Gravel pit operation, open pit mining, strip mining				
Mining (other than gravel, open pit, and strip mining), abandoned mines				
Resource extraction (oil and gas wells and surrounding footprint)				
Agriculture – tilled crop production				
Agriculture – permanent crop (hay pasture, vineyard, orchard, tree plantation)				
Recent old fields and other lands dominated by non-native species (weeds or hay fields)				
Intensively managed golf courses, sports fields, urban parks, expansive lawns				
Vegetation conversion (chaining, cabling, rotochopping, or clear-cutting of woody veg)				
Heavy grazing: (> 2/3 of herbaceous plants have been grazed) by livestock or native ungulate				
Moderate Grazing: (at least 1/3 to 2/3 of herbaceous plants have been grazed) by livestock or native ungulate				
Light Grazing: (< 1/3 of herbaceous plants have been grazed) by livestock or native ungulates				
Heavy browse (> 2/3 of woody plants have been browsed by livestock or native ungulates)				
Moderate browse (at least 1/3 to 2/3 of woody plants have been browsed by livestock or native ungulates)				
Light browse (< 1/3 of woody plants have been browsed by livestock or native ungulates)				
Heavy recreation or human visitation (ATV use / camping / popular fishing spot, etc.)				
Moderate recreation or human visitation (high-use trail)				
Light recreation or human visitation (low-use trail)				
Logging or tree removal with 50-75% of trees				
Selective logging or tree removal with <50% of trees				
Evidence of recent fire (<5years old, still very apparent on vegetation, little regrowth)				
Dam sites and flood disturbed shorelines around water storage reservoirs				
Beetle-killed conifers				
Irrigation ditches, berms, dams, head gates that change how water moves				
Non-native species				
Other:				

Landscape Stressor Comments:

**2. VEGETATION CONDITION METRICS – Check the applicable box.**

**VEGETATION COMPOSITION**



Vegetation compositions and structure, woody regeneration and liter metrics will be calculated out of the field based on the species list and cover values. To aid data interpretation, provide comments on composition and **list noxious species identified in field.**

**2e. REGENERATION OF NATIVE WOODY SPECIES-** Select the statement that best describes the **regeneration of native woody species** within the AA.

Woody species are naturally uncommon or absent.	<b>N/A</b>
All age classes of desirable (native) woody riparian species present.	
Age classes restricted to mature individuals and young sprouts. Middle age groups absent.	
Stand comprised of mainly mature species OR mainly evenly aged young sprouts that choke out other vegetation.	
Woody species predominantly consist of decadent or dying individuals OR woody layer is dominated by Russian olive and/or Salt Cedar	
Regeneration comments and photo #'s:	

**2h. HORIZONTAL INTERSPERSION OF BIOTIC AND ABIOTIC ZONES**

Refer to diagrams below and select the statement that best describes the <b>horizontal interspersion of biotic and abiotic zones</b> within the AA. Rules for defining zones are in the field manual. Include zones of open water when evaluating interspersion.	High degree of horizontal interspersion: AA characterized by a very complex array of nested or interspersed zones with no single dominant zone.	
	Moderate degree of horizontal interspersion: AA characterized by a moderate array of nested or interspersed zones with no single dominant zone.	
	Low degree of horizontal interspersion: AA characterized by a simple array of nested or interspersed zones. One zone may dominate others.	
	No horizontal interspersion: AA characterized by one dominant zone.	



**2k. VEGETATION STRESSORS WITHIN THE AA**

Using the table below, mark all vegetation stressor within the AA. Stressors can overlap (e.g., light grazing can occur along with moderate recreation). Rank the top 3 effecting the wetland.

<i>Vegetation stressor categories</i>	<i>Present</i>	<i>Rank</i>
Unpaved Roads (e.g., driveway, tractor trail, 4-wheel drive roads)		
Vegetation conversion (chaining, cabling, rotochopping, clearcut)		
Logging or tree removal with 50-75% of trees >50 cm dbh removed		
Selective logging or tree removal with <50% of trees >50 cm dbh removed		
Heavy grazing: (> 2/3 of herbaceous plants have been grazed) by livestock or native ungulate		
Moderate Grazing: (at least 1/3 to 2/3 of herbaceous plants have been grazed) by livestock or native ungulate		
Light Grazing: (< 1/3 of herbaceous plants have been grazed) by livestock or native ungulates		
Heavy browse (> 2/3 of woody plants have been browsed) by livestock or native ungulates		
Moderate browse (at least 1/3 to 2/3 of woody plants have been browsed) by livestock or native ungulates		
Light browse (< 1/3 of woody plants have been browsed) by livestock or native ungulates		
Intense recreation or human visitation (ATV use / camping / popular fishing spot, etc.)		
Moderate recreation or human visitation (high-use trail)		
Light recreation or human visitation (low-use trail)		
Recent old fields and other lands dominated by <i>non-native</i> species (weeds or hay)		
Haying of <i>native</i> grassland ( <i>not</i> dominated by non-native hay grasses)		
Beetle-killed conifers		
Non-native Species		
Litter is extensive and limits new growth (thick cattails litter)		
Other:		
Vegetation stressor comments and photo #'s:		

**3. HYDROLOGY METRICS – Check the applicable box.****3a. Water source and Hydrologic stressors within the drainage basin**

Check off all <i>major</i> water sources in the table to the right. If the dominant water source is evident, mark it with a star (*).	<input type="checkbox"/> Overbank flooding	<input type="checkbox"/> Irrigation via direct application
	<input type="checkbox"/> Alluvial aquifer	<input type="checkbox"/> Irrigation via seepage
	<input type="checkbox"/> Groundwater discharge	<input type="checkbox"/> Irrigation via tail water run-off
	<input type="checkbox"/> Natural surface flow	<input type="checkbox"/> Urban run-off / culverts
	<input type="checkbox"/> Precipitation	<input type="checkbox"/> Pipes (directly feeding wetland)
	<input type="checkbox"/> Snowmelt	<input type="checkbox"/> Other:

In the table below, estimate the scope of each **hydrology stressor within the AA and within the 500 m envelope of the AA**. If known hydrologic alterations occur further than 500 m from the AA and are positioned in a way that have an effect on the sites hydrology record the stressors scope in the proper location and please explain in comments below. **Scope rating: 1 = 1–10%, 2 = >10–25%, 3 = >25–50%, 4 = >50–75%, 5 = >75%.**

<i>Hydrology stressor categories</i>	<i>Within AA</i>	<i>Upstream / Upslope</i>	<i>Downstream / Downslope</i>
Ditches < 1 feet deep are present			
Ditches 1 foot to 3 feet deep are present			
Ditches > 3 feet deep are present			
Diversion structures < 1 foot tall are present			

Diversion structures 1 foot to 3 feet tall are present			
Diversion structures > 3 feet tall are present			
Major irrigation canal			
Spring box diverting water from wetland			
Berms present that impede forward or lateral movement of water			
Weir or drop structure that impounds water and controls energy of flow			
Impoundment / stock pond			
Large dam / reservoir			
Dirt or gravel road that alters forward or lateral movement of water			
2-lane road crosses that alters forward or lateral movement of water			
4-lane road crosses that alters forward or lateral movement of water			
Culvert too small to accommodate base flow			
Culvert appears large enough to accommodate base flow but not flood flows			
Culvert appears large enough to accommodate base flow and flood flows			
Pugging by livestock, native ungulates, or wild horses that alters water movement in the site			
Dug pits for holding water			
Fill that has been added to site			
Surrounding land cover / vegetation that interrupts surface flow			
Observed or potential agricultural runoff			
Developed or irrigated lands occupy drainage basin.			
Other:			
Other			
Hydrologic stressor and water source comments:			
<b>Hydrologic landscape and management context. Check all that apply checklist</b>			
Wetland appears to be still connected to its natural water source, natural flows appear to be unaltered.			
Wetland appears to naturally lack water at times.			
Land use in the local watershed is primarily open space or low-density development			
Local watershed includes little or no irrigated land.			
Wetland is in a location that appears to have supported a wetland before development in the immediate drainage basin			
Filling and drawdown of the wetland appear to be unmanaged			
Filling & drawdown are managed to mimic natural timing and amount			
Filling & drawdown are managed with no regard to natural timing and amount			
Xeric vegetation is encroaching into the wetland			
Natural wet-season or dry season inflows to the wetland have been eliminated by impoundment or diversion.			
Wetland exists in intermittent drainage basin that has been bermed or dugout to hold water for livestock use or irrigation storage			
Wetland appears to be largely or entirely supported by anthropogenic inputs such as: direct irrigation, runoff from irrigated fields, seepage from irrigation canals or ditches, urban stormwater runoff, direct pumping, or landscape modification for water storage			

Wetland landscape and management context comments:

#### 4a. WATER SOURCES / INPUTS

Select the statement below that best describes the water sources feeding the AA during the growing season. Use the water source, hydrologic stressor and wetland landscape and management context tables to inform your answers

Water sources are precipitation, groundwater, natural runoff, or natural flow from an adjacent freshwater body. The system may naturally lack water at times, such as in the growing season. There is no indication of direct artificial water sources, either point sources or non-point sources. Land use in the local watershed is primarily open space or low density, passive use with little irrigation.

Water sources are mostly natural, but also include occasional or small amounts of inflow from anthropogenic sources. Indications of anthropogenic sources include developed land or irrigated agriculture that comprises < 20% of the immediate drainage basin, the presence of a few small storm drains or scattered homes with septic system. No large point sources control the overall hydrology.

Water sources are moderately impacted by anthropogenic sources, but are still a mix of natural and non-natural sources. Indications of moderate contribution from anthropogenic sources include developed land or irrigated agriculture that comprises 20–60% of the immediate drainage basin or the presence of a many small storm drains or a few large ones. The key factor to consider is whether the wetland is located in a landscape position that supported a wetland before development and whether the wetland is still connected to its natural water source (e.g., modified ponds on a floodplain that are still connected to alluvial aquifers, natural stream channels that now receive substantial irrigation return flows).

Water sources are primarily from anthropogenic sources (e.g., urban runoff, direct irrigation, pumped water, artificially impounded water, or another artificial hydrology). Indications of substantial artificial hydrology include developed or irrigated agricultural land that comprises > 60% of the immediate drainage basin of the AA, or the presence of major drainage point source discharges that obviously control the hydrology of the AA. The key factor to consider is whether the wetland is located in a landscape position that likely never supported a wetland prior to human development. The reason the wetland exists is because of direct irrigation, irrigation seepage, irrigation return flows, urban storm water runoff, direct pumping, or landscape modifications for water storage.

Natural sources have been **eliminated** based on the following indicators: impoundment of all wet season inflows, diversions of all dry-season inflows, predominance of xeric vegetation, etc. The wetland is in steady decline and may not be a wetland in the near future.

Water Source/ inputs comments:

#### 4b. HYDROPERIOD

Select the statement below that best describes the **hydroperiod** within the AA (extent and duration of inundation and/or saturation). **Use the water source, hydrologic stressor and wetland landscape and management context tables to determine the overall condition of the hydroperiod.** For some wetlands, this may mean that water is being channelized or diverted away from the wetland. For others, water may be concentrated or increased.

Hydroperiod is characterized by natural patterns of filling or inundation and drying or drawdowns. There are no major hydrologic stressors that impact the natural hydroperiod.

Hydroperiod filling or inundation patterns deviate slightly from natural conditions due to presence of stressors such as: small ditches or diversions; berms or roads at/near grade; minor pugging by livestock; or minor flow additions. Outlets may be slightly constricted. Playas are not significantly pitted or dissected. *If hydrology is artificially controlled*, the management regime closely mimics a natural analogue (it is very unusual for a purely artificial wetland to be rated in this category).

Hydroperiod filling or inundation and drying patterns deviate moderately from natural conditions due to presence of stressors such as: ditches or diversions 1–3 ft. deep; two lane roads; culverts adequate for base stream flow but not flood flow; moderate pugging by livestock that could channelize or divert water; shallow pits within playas; or moderate flow additions. Outlets may be moderately constricted, but flow is still possible. *If hydrology is artificially controlled*, the management regime approaches a natural analogue. Site may be passively managed, meaning that the hydroperiod is still connected to and influenced by natural high flows timed with seasonal water levels.

Hydroperiod filling or inundation and drawdown of the AA deviate substantially from natural conditions from high intensity alterations such as: a 4-lane highway; large dikes impounding water; diversions > 3ft. deep that withdraw a significant portion of flow, deep pits in playas; large amounts of fill; significant artificial groundwater pumping; or heavy flow additions. Outlets may be significantly constricted, blocking most flow. <i>If hydrology is artificially controlled</i> , the site is actively managed and not connected to any natural season fluctuations, but the hydroperiod supports natural functioning of the wetland.			
Hydroperiod is dramatically different from natural. Upstream diversions severely stress the wetland. Riverine wetlands may run dry during critical times. <i>If hydrology is artificially controlled</i> , hydroperiod does not mimic natural seasonality. Site is actively managed for filling or drawing down without regard for natural wetland functioning.			
<b>4c. HYDROLOGIC CONNECTIVITY</b>			
Select the statement below that best describes the degree to which <b>hydrology within the AA is connected to the larger landscape</b> throughout the year, but particularly at times of high water. <b>Use the water source, hydrologic stressor and wetland landscape and management context tables to determine the overall condition of hydrologic connectivity.</b> Consider the effect of impoundments, entrenchment, or other obstructions to connectivity that occur within the surrounding landscape, if those impoundments clearly impact the AA.			
<b>General criteria</b>	<b>Riverine variant</b>	<b>Playa variant</b>	
Nothing obstructs lateral or vertical movement of surface or ground water if wetland depends on perched water table then impermeable soil layer (fragipan or duripan) is intact. Rising water in the site has unrestricted access to adjacent upland, without levees, excessively high banks, artificial barriers, or other obstructions to the lateral movement of flood flows.	Completely connected to floodplain (backwater sloughs and channels). No geomorphic modifications have been made to contemporary floodplain. Channel is not entrenched.	Surrounding land cover / vegetation does not interrupt surface flow. No artificial channels feed water to playa.	
Constructed levees or road grades limit the amount of adjacent transition zone or the lateral movement of floodwaters for <50% of the AA boundary. Restrictions may be intermittent along the margins of the AA, or they may occur only along one bank or shore.	Minimally disconnected from floodplain. Up to 25% of stream banks are affected by constructed levees or road grades and/or channel is somewhat entrenched.	Surrounding land cover / vegetation does not interrupt surface flow. Artificial channels may feed minor amounts of water to playa.	
Constructed features such as levees or road grades border 50–90% of the boundary of the AA. Flood flows may overtop the obstructions, but drainage out of the AA is probably obstructed.	Dikes, tide gates, or elevated culverts affect 25-75% of stream banks. Channel may be moderately entrenched and disconnected from the floodplain except in large floods.	Surrounding land cover / vegetation may interrupt surface flow. Artificial channels may feed moderate amounts of excess water to playa.	
Constructed features such as levees or roadbeds border >90% of the boundary of the AA.	Channel is severely entrenched and entirely disconnected from the floodplain.	Surrounding land cover / vegetation may dramatically restrict surface flow. Artificial channels may feed significant amounts of excess water to playa.	
Hydroperiod and hydrologic connectivity comments:			

#### 4. PHYSIOCHEMICAL METRICS – Check the applicable box.

<b>3a. WATER QUALITY - SURFACE WATER TURBIDITY / POLLUTANTS</b>	
Select the statement that best describes the <b>turbidity or evidence or pollutants</b> in surface water within the AA.	
No open water in AA	
No visual evidence of degraded water quality. No visual evidence of turbidity or other pollutants.	
Some negative water quality indicators are present, but limited to small and localized areas within the wetland. Water is slightly cloudy, but there is no obvious source of sedimentation or other pollutants.	
Water is cloudy or has unnatural oil sheen, but the bottom is still visible. Sources of water quality degradation are apparent (identify in comments below). <i>Note: If the sheen breaks apart when you run your finger through it, it is a natural bacterial process and not water pollution. Riverine wetlands can be turbid if flood waters are high</i>	
Water is milky and/or muddy or has unnatural oil sheen. The bottom is difficult to see. There are obvious sources of water quality degradation (identify in comments below). <i>Note: If the sheen breaks apart when you run your finger through it, it is a natural bacterial process and not water pollution. Riverine wetlands can be turbid if flood waters are high</i>	

Surface water turbidity / pollutants comments and photo #'s:

**3b. WATER QUALITY - ALGAL GROWTH**Select the statement that best describes **algal growth** within surface water in the AA.

No open water in AA or evidence of open water.

Water is clear with minimal algal growth.

Algal growth is limited to small and localized areas of the wetland. Water may have a greenish tint or cloudiness.

Algal growth occurs in moderate to large patches throughout the AA. Water may have a moderate greenish tint or sheen. Sources of water quality degradation are apparent (identify in comments below).

Algal mats are extensive, blocking light to the bottom. Water may have a strong greenish tint and the bottom is difficult to see. There are obvious sources of water quality degradation (identify in comments below).

Algal growth comments and photo #'s:

*Algal growth may be natural and not necessarily indicative of poor water quality. If algal growth appears natural, describe and record % of total algae that is due to natural processes.***3c. SUBSTRATE / SOIL DISTURBANCE**

Select the statement below that best describes disturbance to the substrate or soil within the AA. For playas, the most significant substrate disturbance is sedimentation or unnaturally filling, which prevents the system's ability to pond after heavy rains. For other wetland types, disturbances may lead to bare or exposed soil and may increase ponding or channelization where it is not normally. For any wetland type, consider the disturbance relative to what is expected for the system.

No soil disturbance within AA. Little bare soil OR bare soil areas are limited to naturally caused disturbances such as flood deposition or game trails OR soil is naturally bare (e.g., playas). No pugging, soil compaction, or sedimentation.

Less than 10% of the AA affected by some amount of bare soil, pugging, compaction, or sedimentation present due to human causes. The depth of disturbance is limited to 1 – 2 inches and does not show evidence of altering hydrology or vegetation growth at the site

10 –25% of the AA has bare soil areas due to human causes are common. There may be pugging due to livestock resulting in several inches of soil disturbance. Sedimentation may be filling the wetland. Damage is obvious, but not excessive.

25-50% of the AA has bare soil areas due to human causes are common. ORVs or other machinery may have left some shallow ruts &lt; 3 inches deep or livestock pugging and/or trails are widespread. Unnatural hummocks created by livestock, wild horses, or native ungulates present, especially when the site lacks hummock forming vegetation. These hummocks typically have sheer edges with exposed soil. Compaction and disturbance change water moment in the site and affect vegetation growth. Sedimentation may have severely impacted the hydrology.

Greater than 50% off the AA has bare soil areas that substantially degrade the site and have led to altered hydrology or other long-lasting impacts. Deep ruts from ORVs or machinery may be present, Unnatural hummocks created by livestock, wild horses, or native ungulates present, especially when the site lacks hummock forming vegetation. These hummocks typically have sheer edges with exposed soil. Sedimentation has dried the wetland.

Substrate / soil comments and photo #'s:

**3d. PHYSIOCHEMICAL STRESSORS WITHIN THE AA**Using the table below, estimate the independent scope of each physiochemical stressor within the AA. Independent scopes can overlap (e.g., soil compaction can occur with trash or refuse). **Scope rating: 1 = 1–10%, 2 = >10–25%, 3 = >25–50%, 4 = >50–75%, 5 = >75%.***Physiochemical stressor categories**Scope*

Erosion

Sedimentation

Current plowing or disking

Historic plowing or disking (evident by abrupt A horizon boundary at plow depth)

Substrate removal (excavation)

Filling or dumping of sediment

Trash or refuse dumping

Compaction and soil disturbance by livestock, wild horses, or native ungulates &lt; 3 inches deep

Compaction and soil disturbance by livestock, wild horses, or native ungulates &gt; 3 inches deep

Unnatural hummocks created by livestock, wild horses, or native ungulates. These typically have sheer edges with exposed soil. Site lacks hummock forming vegetation

Compaction and soil disturbance by human use (trails, ORV use, camping) < 3 inches deep	
Compaction and soil disturbance by human use (trails, ORV use, camping) > 3 inches deep	
Mining activities, current or historic	
Obvious point source of water pollutants (discharge from waste water plants, factories)	
Agricultural runoff (drain tiles, excess irrigation)	
Direct application of agricultural chemicals	
Discharge or runoff from feedlots	
Obvious excess salinity (dead or stressed plants, salt encrustations)	
Other:	
Physiochemical stressor comments:	



# AREM Long Form

Please evaluate the wetland or riparian habitat within the **200 meter** buffer when answering the below questions. Do not consider upland habitat except for questions 16 – 21. For each numbered item, check only one response unless noted otherwise. Then proceed to the next question unless noted otherwise. Parenthetical names are the names of fields in the supporting software database (WHRBASE). If a field name is lacking, the information is not used directly.

**Note: 1 Acre = .5 hectares**

1. Season: Migratory \_\_\_\_\_ Breeding \_\_\_\_\_ Winter \_\_\_\_\_
2. LOCATION. Is the area part of, or is it within 0.5 mile of, a major\* river or lake?  
*\* river channel wider than 100 ft, or lake larger than 40 acres*  
 \_\_\_\_ Yes (field BigWater) \_\_\_\_ No
3. SURFACE WATER. During this season, does the area contain at least 0.1 acre\* of surface water, either obscured by vegetation or not?  
*\* See Figure B-1 for guidance in estimating acreage categories.*  
 \_\_\_\_ Yes (field AnyWater). Go to next question.  
 \_\_\_\_ No. **Skip to question #5.**
4. OPEN WATER. During this season, how much open\* water is present in the area?  
*\* water deeper than 2 inches and mostly lacking vegetation (except submerged plants).*  
 \_\_\_\_ > 20 acres and it is mostly wider than 500 ft (field OpenBig)  
 \_\_\_\_ < 1 acre, or, >1 acre but mostly narrower than 3 ft (field OpenSmall)  
 \_\_\_\_ Other conditions (field OpenOther)
5. SPECIFIC AQUATIC CONDITIONS  
 Check all that apply during this season:  
 \_\_\_\_ > 0.1 acre of the surface water is still, i.e., usually flows at less than 1 ft/s (field StillWater)  
 \_\_\_\_ The evaluated area can be assumed to contain fish (field Fish)  
 \_\_\_\_ The evaluated area can be assumed to contain frogs, salamanders, and/or crayfish (field Amphibs)  
 \_\_\_\_ Water transparency in the deepest part of the area is (or would be, if depth is shallow) sufficient to see an object 10 inches below the surface, and the area is not known to have problems with metal contamination (field Clear)  
 \_\_\_\_ The evaluated area is highly enriched by direct fertilizer applications, water from nearby feedlots, or other sources (field Enriched)  
 \_\_\_\_ Most of the normally-flooded part of the area goes dry at least one year in five, or, is subject to flooding from a river at least as often (field Drawdown)
6. BARE SOIL. Is there at least 0.1 acre of mud\*, alkali flat, gravel/sand bar, recently tilled soil, and/or heavily grazed open (grassy, non-shrubby) areas during this season?  
*\* includes soil that is continually saturated up to the surface, or which was previously covered by water but has become exposed to the air during this period*  
 \_\_\_\_ Yes (field Bare). Go to next question.  
 \_\_\_\_ No. **Skip to question #7.**
7. LARGE MUDFLAT. Does the area at this season contain mud that has all these features?:
  - At least 1 acre in size
  - Maximum dimension is greater than 100 ft
  - Salt crust or salt stains are not apparent
  - Not recessed within a wash or canal whose depth (relative to surrounding landscape) is greater than half its width.  
 \_\_\_\_ Yes (field MudBig) \_\_\_\_ No

8. TREES. Are there at least 3 trees\*:  
*\* woody plants taller than 20 ft.*  
 \_\_\_\_ in the evaluation area? (field TreeIn).  
 \_\_\_\_ within 1000 ft of the evaluation area? (field TreeNear). **Go to #8.**  
 \_\_\_\_ neither of the above. **Skip to #11.**
9. TREE COVER. Check one or more responses below that describe the maximum cumulative acreage of various conditions of tree cover in the evaluation area. Also include areas within 300 ft:  
 \_\_\_\_ >1 acre, dense\*, and wide\*\* (field ForestDens)  
 \_\_\_\_ >1 acre and open; or, dense but narrow (field ForestOpen)  
 \_\_\_\_ 0.1–1 acre, dense\* (field WoodDens)  
 \_\_\_\_ 0.1–1 acre, open (field WoodOpen)  
 \_\_\_\_ <0.1 acre  
*\* Dense= the tree canopy, viewed from the ground during midsummer, appears at least 50% closed, as averaged across an area that is at least as large as the acreage specified.*  
*\*\* Wide= the wooded area is wider than 300 ft (average).*
10. BIG TREES. Are there at least three trees whose trunk diameter 20 ft above the ground is >12 inches?  
 \_\_\_\_ Yes (field TreesBig) \_\_\_\_ No
11. SNAGS. Are there at least three snags, or trees with dead limbs with diameter >5 inches?  
 \_\_\_\_ Yes (field Snags) \_\_\_\_ No
12. SHRUBS. Is there at least 0.1 acre of shrubs\*:  
*\* woody plants 2–20 ft in height.*  
 \_\_\_\_ in the evaluation area? (field ShrubIn).  
 \_\_\_\_ within 1000 ft of the wetland (including the wetland itself)? (field ShrubNear). **Go to #12.**  
 \_\_\_\_ Neither of the above. **Skip to #13.**
13. SHRUB SPECIES AND DENSITY. Check one or more responses below that describe the maximum cumulative extent of various types and conditions of shrub cover in the evaluation area. Also include areas within 300 ft.  
 Willow:  
 \_\_\_\_ >1 acre, dense\*, and wide\*\* (field WwMuchDens)  
 \_\_\_\_ >1 acre and open; or, dense but narrow (field WwMuchOpen)  
 \_\_\_\_ 0.1–1 acre, dense\* (field WwSomeDens)  
 \_\_\_\_ 0.1–1 acre, open (field WwSomeOpen)  
 \_\_\_\_ <0.1 acre; or larger area but height mostly <4 ft and openly spaced
- Greasewood or other tall desert shrubs:  
 \_\_\_\_ >1 acre, dense\*, and wide\*\* (field GrMuchDens)  
 \_\_\_\_ >1 acre and open; or, dense but narrow (field GrMuchOpen)  
 \_\_\_\_ 0.1–1 acre, dense\* (field GrSomeDens)  
 \_\_\_\_ 0.1–1 acre, open (field GrSomeOpen)  
 \_\_\_\_ <0.1 acre
- Russian olive, sumac, buffaloberry, wild rose, or others with fleshy fruit:  
 \_\_\_\_ >1 acre, dense\*, and wide\*\* (field FrMuchDens)  
 \_\_\_\_ >1 acre, open; or, dense but narrow (field FrMuchOpen)  
 \_\_\_\_ 0.1–1 acre, dense (field FrSomeDens)  
 \_\_\_\_ 0.1–1 acre, open (field FrSomeOpen)  
 \_\_\_\_ <0.1 acre; or larger area but height mostly <4 ft

Tamarisk (salt cedar):

- \_\_\_\_\_ >1 acre, dense\*, and wide\*\* (field TmMuchDens)
- \_\_\_\_\_ >1 acre, open; or, dense but narrow (field TmMuchOpen)
- \_\_\_\_\_ 0.1–1 acre, dense (field TmSomeDens)
- \_\_\_\_\_ 0.1–1 acre, open (field TmSomeOpen)
- \_\_\_\_\_ <0.1 acre; or larger area but height mostly <4 ft

\* *Dense*= the shrub canopy, as viewed from a height of 100 ft during midsummer, appears to be >50% closed, as averaged across an area that is at least as large as the acreage specified.

\*\* *Wide*= the shrub area is wider than 300 ft (average).

14. HERBACEOUS VEGETATION. Is there at least 0.1 acre of herbaceous vegetation\*:

\* *Nonwoody plants such as cattail, bulrush, sedges, grasses, and forbs.*

- \_\_\_\_\_ in the evaluation area? (field HerbIn).
- \_\_\_\_\_ within 1000 ft? (field HerbNear). **Go to #14.**
- \_\_\_\_\_ Neither of the above. **Skip to #15.**

15. HERBACEOUS SPECIES. Check one or more responses below that describe the maximum cumulative extent of various types and conditions of shrub cover in the evaluation area. Also include areas within 300 ft.

Robust emergents (e.g., cattail, phragmites)

- \_\_\_\_\_ >1 acre, dense\*, and wide\*\* (field RbMuchDens)
- \_\_\_\_\_ >1 acre, open; or dense but narrow (field RbMuchOpen)
- \_\_\_\_\_ 0.1–1 acre, dense (field RbSomeDens)
- \_\_\_\_\_ 0.1–1 acre, open (field RbSomeOpen)

Other wet\*\* emergents (e.g., bulrush, sedge)

- \_\_\_\_\_ >1 acre, dense\*, wide\*\*, and tall\*\*\* (field WEMuchDens)
- \_\_\_\_\_ >1 acre, tall, open; or dense but narrow (field WEMuchOpen)
- \_\_\_\_\_ >1 acre, dense or open, and short (field WEMuchShrt)
- \_\_\_\_\_ 0.1–1 acre, tall, dense (field WESomeDens)
- \_\_\_\_\_ 0.1–1 acre, tall, open; or dense but narrow (field WESomeOpen)
- \_\_\_\_\_ 0.1–1 acre, dense or open, and short (field WESomeShrt)

Drier emergents (e.g., saltgrass, other grasses)

- \_\_\_\_\_ >1 acre, dense\*, wide\*\*, and tall\*\*\* (field DEMuchDens)
- \_\_\_\_\_ >1 acre, tall, open; or dense but narrow (field DEMuchOpen)
- \_\_\_\_\_ >1 acre, dense or open, and short (field DEMuchShrt)
- \_\_\_\_\_ 0.1–1 acre, tall, dense (field DESomeDens)
- \_\_\_\_\_ 0.1–1 acre, tall, open; or dense but narrow (field DESomeOpen)
- \_\_\_\_\_ 0.1–1 acre, dense or open, and short (field DESomeShrt)

Broad-leaved Forbs (e.g., milkweed, thistle, alfalfa)

- \_\_\_\_\_ >1 acre (field ForbMuch)
- \_\_\_\_\_ 0.1–1 acre (field ForbSome)

Aquatic plants (e.g., watercress, sago pondweed, duckweed)

- \_\_\_\_\_ >10 acres (field AqMuch)
- \_\_\_\_\_ 0.1–10 acres (field AqSome)

\* *Dense*= plants are so close together that the duff layer or soil beneath the plants is mostly obscured by foliage, when looking down from just above the plant tops.

\*\* *Wet*= water is visible at or above the soil surface during most of the growing season.

\*\*\* *Wide*= the shrub area is wider than 300 ft (average).

\*\*\*\* *Tall*= taller than 1 ft.

16. SURROUNDING LAND COVER (includes wetland and upland habitat). Check one:

Within 0.5 mi of the wetland, >60% of the land cover is:

\_\_\_\_ Pasture, alfalfa, grain crops, row crops, other wetlands, grass lawns, and/or weed fields (field SurAgwet)

\_\_\_\_ Desert shrubs (e.g., sagebrush, shadscale, rabbitbrush)(field SurDesrt)

\_\_\_\_ Pinyon-juniper (field SurPJ)

\_\_\_\_ Oak scrub (e.g., Gambel oak, serviceberry, skunkbrush)(field SurOak)

\_\_\_\_ Other, or none of the above comprise >60%

17. LOCAL LAND COVER (includes wetland and upland habitat). Check one:

Within 3 mi of the wetland, > 60% of the land cover is:

\_\_\_\_ Pasture, alfalfa, grain crops, row crops, other wetlands, grass lawns, and/or weed fields (field LocAgWet)

\_\_\_\_ Desert shrubs (e.g., sagebrush, shadscale, rabbitbrush)(field LocDesrt)

\_\_\_\_ Pinyon-juniper (field LocPJ)

\_\_\_\_ Oak scrub (e.g., Gambel oak, serviceberry, skunkbrush)(field LocOak)

\_\_\_\_ Other, or none of the above comprise >60%

18. VISUAL SECLUSION

Check only one:

\_\_\_\_ Both of the following:

(a) wetland is seldom visited by people on foot or boat (less than once weekly), (b) there are no paved roads within 600 ft, or if there are, wetland is not visible from the roads (field SeclusionH).

\_\_\_\_ Either (a) or (b) above (field SeclusionM).

\_\_\_\_ Other condition.

19. PREDATION POTENTIAL

Check only one. The evaluation area:

\_\_\_\_ is linear\*, adjoins a heavily-traveled road (usual maximum of >1 car/minute), and/or is in a high-density housing area (>1 house/5 acres) (field PredHPot)

\_\_\_\_ adjoins a less-traveled road, and/or is in an area with sparser housing density but is closer than 1000 ft to a normally-occupied building (field PredMPot)

\_\_\_\_ Other condition.

\* at least 90% of the area being evaluated is within 25 ft of a canal, road, railroad tracks, or other artificially linear feature.

20. GRAZED, BURNED, MOWED. Is the area mowed, burned, or grazed intensively (i.e., with clearly visible effects on vegetation) during this season?

\_\_\_\_ Yes (field GrazBurnMo)

\_\_\_\_ No

21. NESTING LOCATIONS

Check all that apply:

\_\_\_\_ Semi-open structures (bridges, barns) suitable for nesting swallows are present within 300 ft (field SwallNest)

\_\_\_\_ Platforms suitable for nesting geese are present in the wetland or along its perimeter (field GooseNest)

\_\_\_\_ Vertical, mostly bare dirt banks at least 5 ft high are present within 0.5 mi., of potential use to nesting kingfishers, barn owls, and swallows (field Banks)

**APPENDIX D. Scoring formulas for Ecological Integrity Assessment wetland condition scores.**

**Table D.1.** EIA ranks and definitions adapted from (Lemly and Gilligan 2013).

<i>Rank</i>	<i>Condition Category</i>	<i>Interpretation</i>
<b>A</b>	<b>Excellent / Reference Condition (No or Minimal Human Impact)</b>	Wetland functions within the bounds of natural disturbance regimes. The surrounding landscape contains natural habitats that are essentially unfragmented with little to no stressors; vegetation structure and composition are within the natural range of variation, nonnative species are essentially absent, and a comprehensive set of key species are present; soil properties and hydrological functions are intact. Management should focus on preservation and protection.
<b>B</b>	<b>Good / Slight Deviation from Reference</b>	Wetland predominantly functions within the bounds of natural disturbance regimes. The surrounding landscape contains largely natural habitats that are minimally fragmented with few stressors; vegetation structure and composition deviate slightly from the natural range of variation, nonnative species and noxious weeds are present in minor amounts, and most key species are present; soils properties and hydrology are only slightly altered. Management should focus on the prevention of further alteration.
<b>C</b>	<b>Fair / Moderate Deviation from Reference</b>	Wetland has a number of unfavorable characteristics. The surrounding landscape is moderately fragmented with several stressors; the vegetation structure and composition is somewhat outside the natural range of variation, nonnative species and noxious weeds may have a sizeable presence or moderately negative impacts, and many key species are absent; soil properties and hydrology are altered. Management would be needed to maintain or restore certain ecological attributes.
<b>D</b>	<b>Poor / Significant Deviation from Reference</b>	Wetland has severely altered characteristics. The surrounding landscape contains little natural habitat and is very fragmented; the vegetation structure and composition are well beyond their natural range of variation, nonnative species and noxious weeds exert a strong negative impact, and most key species are absent; soil properties and hydrology are severely altered. There may be little long term conservation value without restoration, and such restoration may be difficult or uncertain.

**Table D.2. EIA methods for scoring.**

1. The score for each EIA submetric was calculated using the equations below.

Landscape Context Score:

$$(\text{Landscape Fragmentation} * 0.4) + ([(\text{Buffer Width} * \text{Buffer Extent})^{1/2} * ((\text{Buffer Condition} + \text{Buffer Natural Cover})/2)]^{1/2} * 0.6)$$

Biotic Condition Score:

$$(\text{Relative Cover Native Plant Sp.} * 0.2) + (\text{Absolute Cover Noxious Weeds} * 0.2) + (\text{Mean C} * 0.4) + (\text{Horizontal Interspersion} * 0.2)$$

Hydrologic Condition Score:

Landscape Hydrology Metric score

Physicochemical Condition Score:

$$(\text{Surface Water Quality} * 0.25) + (\text{Algal Growth} * 0.25) + (\text{Substrate/Soil Disturbance} * 0.5)$$

*If no standing water was present, score = Substrate/Soil Disturbance.*

2. EIA score was calculated using submetric scores:

EIA Score:

$$(\text{Landscape Context} * 0.2) + (\text{Biotic Condition} * 0.4) + (\text{Hydrologic Condition} * 0.3) + (\text{Physicochemical Condition} * 0.1)$$

3. Score to rank conversion:

A = 4.5 – 5.0

B = 3.5 – <4.5

C = 2.5 - <3.5

D = 1.0 - <2.5

## Appendix E: Indicators of Disturbance Categories

Category	Stressor Type	Landscape	Hydrology	Physiochemistry	Vegetation
<b>Agriculture</b>	Agriculture – permanent crop (hay pasture, vineyard, orchard, tree plantation)	x			
	Agriculture – tilled crop production	x			
	Current plowing or disking			x	
	Haying of native grassland (not dominated by non-native hay grasses)				x
	Recent old fields and other fallow lands dominated by non-native species (weeds or hay)	x			x
<b>Browse</b>	Heavy browse (at least 2/3 of woody plants have been browsed) by livestock or native ungulates	x			x
	Moderate browse (at least 1/3 to 2/3 of woody plants have been browsed) by livestock or native ungulates	x			x
	Light browse (< 1/3 of woody plants have been browsed) by livestock or native ungulates	x			x
<b>Development</b>	Domestic or commercially developed buildings	x			
<b>Filling</b>	Filling or dumping of sediment		x	x	
	Sedimentation			x	
<b>Grazing</b>	Heavy grazing: (> 2/3 of herbaceous plants have been grazed) by livestock or native ungulate	x			x
	Moderate Grazing: (at least 1/3 to 2/3 of herbaceous plants have been grazed) by livestock or native ungulate	x			x
	Light Grazing: (< 1/3 of herbaceous plants have been grazed) by livestock or native ungulates	x			x

Category	Stressor Type	Landscape	Hydrology	Physiochemistry	Vegetation
<b>Invasive Species</b>	Invasive Species	x			x
<b>Irrigation Infrastructure</b>	Ditches < x feet deep are present		x		
	Ditches > 3 feet deep are present		x		
	Ditches x foot to 3 feet deep are present		x		
	Diversion structures < x foot tall are present		x		
	Diversion structures > 3 feet tall are present		x		
	Diversion structures x foot to 3 feet tall are present		x		
	Irrigation ditches, berms, head gates that change how water moves	x			
	Spring box diverting water from wetland		x		
	Weir or drop structure that impounds water and controls energy of flow		x		
<b>Natural</b>	Beetle-killed conifers	x			x
	Evidence of recent fire (<5 years old, still very apparent on vegetation, little regrowth)	x			x
<b>Pollution</b>	Agricultural runoff (drain tiles, excess irrigation)			x	
	Direct application of agricultural chemicals			x	
	Discharge or runoff from feedlots			x	
	Observed or potential agricultural runoff		x		
	Obvious point source of water pollutants (discharge from waste water plants, factories)			x	
	Trash or refuse dumping	x		x	



Category	Stressor Type	Landscape	Hydrology	Physiochemistry	Vegetation
<b>Recreation</b>	Intense recreation or human visitation (ATV use / camping / popular fishing spot, etc.)	x			x
	Intensively managed golf courses, sports fields, urban parks, expansive lawns	x			
	Moderate recreation or human visitation (high-use trail)	x			x
	Light recreation or human visitation (low-use trail)	x			x
<b>Resource Extraction</b>	Gravel pit operation, open pit mining, strip mining	x			
	Logging or tree removal with 50-75% of trees >50 cm dbh removed	x			
	Mining (other than gravel, open pit, and strip mining), abandoned mines	x			
	Resource extraction (oil and gas wells and surrounding footprint)	x			
<b>Roads</b>	2-track or lightly used farm road that coauses fragmentation or alters the lateral movement of water	x	x		x
	2-lane road crosses that alters forward or lateral movement of water		x		
	4-lane road crosses that alters forward or lateral movement of water		x		
	Culvert appears large enough to accommodate base flow and flood flows		x		
	Culvert appears large enough to accommodate base flow but not flood flows		x		
	Culvert too small to accommodate base flow		x		
	Paved roads, parking lots, railroad tracks	x			

Category	Stressor Type	Landscape	Hydrology	Physiochemistry	Vegetation
<b>Soil Degradation</b>	Compaction and soil disturbance by human use (trails, ORV use, camping) < 3 inches deep			x	
	Compaction and soil disturbance by human use (trails, ORV use, camping) > 3 inches deep			x	
	Compaction and soil disturbance by livestock, wild horses, or native ungulates < 3 inches deep			x	
	Compaction and soil disturbance by livestock, wild horses, or native ungulates > 3 inches deep			x	
	Erosion			x	
	Historic plowing or disking (evident by abrupt A horizon boundary at plow depth)			x	
	Mining activities, current or historic			x	
	Obvious excess salinity (dead or stressed plants, salt encrustations)			x	
	Pugging by livestock, native ungulates, or wild horses that alters water movement in the site		x	x	
	Substrate removal (excavation)			x	
<b>Veg Conversion</b>	Litter is extensive and limits new growth (thick cattails litter)				x
	Logging or tree removal with 50-75% of trees >50 cm dbh removed				x
	Selective logging or tree removal with <50% of trees >50 cm dbh removed	x			x
	Surrounding land cover / vegetation that interrupts surface flow		x		
	Vegetation conversion (chaining, cabling, rotochopping, clear-cut)	x			x
<b>Water Development</b>	Berms present that impede forward or lateral movement of water		x		
	Dam sites and flood disturbed shorelines around water storage reservoirs	x			
	Dug pits for holding water		x		
	Impoundment / stock pond		x		
	Large dam / reservoir		x		
	Major irrigation canal		x		