

2009 Egyptian Cottonworm (*Spodoptera littoralis*) Survey Summary

Report Generated:	9/16/2009 8:49
Data Range:	5/26/2009 - 8/26/2009

* Generated by Wyoming Pest Detection Program - www.uwyo.edu/capsweb

COUNTY	S. LITTORALIS TRAPS OR SURVEY SITES		NEW COUNTY RECORD
	NEGATIVE	POSITIVE	
ALBANY	0	0	NO
BIG HORN	7	0	NO
CAMPBELL	0	0	NO
CARBON	0	0	NO
CONVERSE	0	0	NO
CROOK	0	0	NO
FREMONT	5	0	NO
GOSHEN	6	0	NO
HOT SPRINGS	3	0	NO
JOHNSON	0	0	NO
LARAMIE	6	0	NO
LINCOLN	0	0	NO
NATRONA	0	0	NO
NIOBRARA	0	0	NO
PARK	7	0	NO
PLATTE	6	0	NO
SHERIDAN	0	0	NO
SUBLETTE	0	0	NO
SWEETWATER	0	0	NO
TETON	0	0	NO
UINTA	0	0	NO
WASHAKIE	5	0	NO
WESTON	0	0	NO
YELLOWSTONE NATIONAL PARK	0	0	NO
TOTAL	45	0	0

S. littoralis was not found in Wyoming in 2009

45 Traps or Survey Sites were examined for S. littoralis in 8 Wyoming Counties

- 0 Traps or Survey Sites were in Apple
- 0 Traps or Survey Sites were in Plum
- 0 Traps or Survey Sites were in Crabapple
- 27 Traps or Survey Sites were in Barley
 - 1 Trap or Survey Site was in Oats
- 17 Traps or Survey Sites were in Wheat

45 Traps were examined:

- 43 Traps were placed in Crops
 - 0 Traps were placed in Home Gardens or Yards
 - 0 Traps were placed in Nurseries
 - 2 Traps were placed in Research / Experiment Areas