

2010 Egyptian Cottonworm (*Spodoptera littoralis*) Survey Summary

Report Generated:	1/21/2011 12:51
Data Range:	5/24/2010 - 9/21/2010

* Generated by Wyoming Pest Detection Program - www.uwyo.edu/capsweb

COUNTY	S. LITTORALIS TRAPS OR SURVEY SITES		NEW COUNTY RECORD
	NEGATIVE	POSITIVE	
ALBANY	2	0	NO
BIG HORN	4	0	NO
CAMPBELL	0	0	NO
CARBON	0	0	NO
CONVERSE	0	0	NO
CROOK	0	0	NO
FREMONT	6	0	NO
GOSHEN	13	0	NO
HOT SPRINGS	3	0	NO
JOHNSON	0	0	NO
LARAMIE	8	0	NO
LINCOLN	0	0	NO
NATRONA	0	0	NO
NIOBRARA	0	0	NO
PARK	5	0	NO
PLATTE	7	0	NO
SHERIDAN	3	0	NO
SUBLETTE	0	0	NO
SWEETWATER	0	0	NO
TETON	0	0	NO
UINTA	0	0	NO
WASHAKIE	8	0	NO
WESTON	0	0	NO
YELLOWSTONE NATIONAL PARK	0	0	NO
TOTAL	59	0	0

S. littoralis was not found in Wyoming in 2010

59 Traps or Survey Sites were examined for S. littoralis in 10 Wyoming Counties

- 2 Traps or Survey Sites were in Apple
- 0 Traps or Survey Sites were in Plum
- 0 Traps or Survey Sites were in Crabapple
- 12 Traps or Survey Sites were in Grape
- 26 Traps or Survey Sites were in Barley
- 0 Traps or Survey Sites were in Oats
- 19 Traps or Survey Sites were in Wheat

59 Traps were examined:

- 44 Traps were placed in Crops
- 0 Traps were placed in Home Gardens or Yards
- 2 Traps were placed in Nurseries
- 4 Traps were placed in Research / Experiment Areas
- 9 Traps were placed in Vineyards