

2017 Egyptian Cottonworm (*Spodoptera littoralis*) Survey Summary

Report Generated:	9/18/2017 11:20
Data Range:	6/5/2017 - 8/25/2017

* Generated by Wyoming Pest Detection Program - www.uwyo.edu/capsweb

COUNTY	S. LITTORALIS TRAPS OR SURVEY SITES		NEW COUNTY RECORD
	NEGATIVE	POSITIVE	
ALBANY	0	0	NO
BIG HORN	5	0	NO
CAMPBELL	0	0	NO
CARBON	0	0	NO
CONVERSE	0	0	NO
CROOK	0	0	NO
FREMONT	0	0	NO
GOSHEN	5	0	NO
HOT SPRINGS	0	0	NO
JOHNSON	0	0	NO
LARAMIE	5	0	NO
LINCOLN	5	0	NO
NATRONA	0	0	NO
NIOBRARA	0	0	NO
PARK	5	0	NO
PLATTE	5	0	NO
SHERIDAN	0	0	NO
SUBLETTE	0	0	NO
SWEETWATER	0	0	NO
TETON	0	0	NO
UINTA	0	0	NO
WASHAKIE	5	0	NO
WESTON	0	0	NO
YELLOWSTONE NATIONAL PARK	0	0	NO
TOTAL	35	0	0

S. littoralis was not found in Wyoming in 2017

35 Traps or Survey Sites were examined for S. littoralis in 7 Wyoming Counties

- 0 Traps or Survey Sites were in Apple
- 0 Traps or Survey Sites were in Plum
- 0 Traps or Survey Sites were in Crabapple
- 0 Traps or Survey Sites were in Grape
- 22 Traps or Survey Sites were in Barley
- 0 Traps or Survey Sites were in Oats
- 13 Traps or Survey Sites were in Wheat

35 Traps were examined:

- 33 Traps were placed in Crops
- 0 Traps were placed in Home Gardens or Yards
- 0 Traps were placed in Nurseries
- 0 Traps were placed in Orchards
- 2 Traps were placed in Research / Experiment Areas
- 0 Traps were placed in Vineyards