2020 Egyptian Cottonworm (Spodoptera littoralis) Survey Summary		
Report Generated:	1/5/2021 12:58	
Data Range:	6/11/2020 - 8/25/2020	

* Generated by Wyoming Pest Detection Program - www.uwyo.edu/capsweb

	S. LITTORALIS TRAPS OR SURVEY SITES		NEW COUNTY
COUNTY	NEGATIVE	POSITIVE	RECORD
ALBANY	0	0	NO
BIG HORN	0	0	NO
CAMPBELL	0	0	NO
CARBON	0	0	NO
CONVERSE	0	0	NO
CROOK	0	0	NO
FREMONT	0	0	NO
GOSHEN	8	0	NO
HOT SPRINGS	0	0	NO
JOHNSON	0	0	NO
LARAMIE	6	0	NO
LINCOLN	0	0	NO
NATRONA	0	0	NO
NIOBRARA	0	0	NO
PARK	0	0	NO
PLATTE	6	0	NO
SHERIDAN	0	0	NO
SUBLETTE	0	0	NO
SWEETWATER	0	0	NO
TETON	0	0	NO
UINTA	0	0	NO
WASHAKIE	0	0	NO
WESTON	0	0	NO
YELLOWSTONE			
NATIONAL PARK	0	0	NO
TOTAL	20	0	0

S. littoralis was not found in Wyoming in 2020

20 Traps or Survey Sites were examined for S. littoralis in 3 Wyoming Counties

- 0 Traps or Survey Sites were in Apple
- 0 Traps or Survey Sites were in Plum
- 3 Traps or Survey Sites were in Corn
- 0 Traps or Survey Sites were in Crabapple
- 3 Traps or Survey Sites were in Grape
- 2 Traps or Survey Sites were in Barley
- 0 Traps or Survey Sites were in Oats
- 1 Trap or Survey Site was in Spelt
- 11 Traps or Survey Sites were in Wheat

20 Traps were examined:

- 16 Traps were placed in Crops
- 0 Traps were placed in Home Gardens or Yards
- 0 Traps were placed in Nurseries
- 0 Traps were placed in Orchards
- 1 Trap was placed in a Research / Experiment Area
- 3 Traps were placed in Vineyards